

昭和基地で観測された磁気嵐時高エネルギー降下電子と VLF 放射の統計的特性

田所裕康<sup>1</sup>、三好由純<sup>2</sup>、山岸久雄<sup>1</sup>、宮岡宏<sup>1</sup>、田中良昌<sup>1</sup>

<sup>1</sup> 国立極地研究所

<sup>2</sup> 名古屋大学太陽地球環境研究所

## Statistical characteristics of storm-time energetic electron precipitation and VLF emissions at Syowa station

Hiroyasu Tadokoro<sup>1</sup>, Yoshizumi Miyoshi<sup>2</sup>, Hisao Yamagishi<sup>1</sup>, Hiroshi Miyaoka<sup>1</sup> and Yoshimasa Tanaka<sup>1</sup>

<sup>1</sup>*National Institute of Polar Research*

<sup>2</sup>*STEL, Nagoya University*

Radiation belt electrons show dynamic variations during magnetic storms. Wave-particle interactions by whistler mode waves, especially chorus waves, are important to understand acceleration and loss processes of the energetic electrons.

To understand wave-particle interactions during the magnetic storms, we have analyzed VLF data at 750 Hz, 1.2kHz, and 2.0kHz from the VLF receiver and Cosmic Noise Absorption (CNA) data from the riometer, which reflects variations of energetic electron precipitation, at Syowa station (L=6.1). Statistical variations of CNA distributions as a function of magnetic local time (MLT) during 43 storms in 2005-2009, with minimum Dst -30nT to -100 nT, show enhanced energetic electron precipitation in the morning side during the recovery phase of these storms. This would be related to drift path of the energetic electrons in magnetosphere. To examine the characteristics of precipitating electrons, we have analyzed VLF data at 750Hz, 1.2kHz, and 2.0kHz. Strong VLF emissions are observed at the morning side during the storm recovery phase. This is almost correlated with the strong energetic electron precipitating region. The characteristics of the VLF emissions and energetic electron precipitation during the storms will be discussed in this presentation.