

# 顕微ラマン分光法による南極ドームふじアイスコアに含まれる 微液胞と微粒子の化学組成

櫻井俊光<sup>1</sup>、大野浩<sup>2</sup>、内田努<sup>3</sup>、本山秀明<sup>1,4</sup>

<sup>1</sup> 国立極地研究所

<sup>2</sup> 北見工業大学

<sup>3</sup> 北海道大学大学院工学研究院

<sup>4</sup> 総合研究大学院大学

## Chemical composition of micro-droplets and solid microparticles in the Dome Fuji deep ice core, Antarctica : using micro-Raman spectroscopy

T. Sakurai<sup>1</sup>, H. Ohno<sup>2</sup>, T. Uchida<sup>3</sup>, H. Motoyama<sup>1,4</sup>

<sup>1</sup> *National Institute of Polar Research*

<sup>2</sup> *Kitami Institute of Technology*

<sup>3</sup> *Graduate School of Engineering, Hokkaido University*

<sup>4</sup> *Department of Polar Science, The Graduate University for Advanced Studies (SOKENDAI)*

Climatic signals in deep ice cores, particularly from ion concentrations, may be affected by the diffusion of liquid solution along grain boundaries of ice. Such solutions include sulfates. Because of the difficulty of detecting sulfate liquids in the ice matrix, we must infer the phase state of the sulfates from the ice temperature and inclusion properties. In this study, we use micro-Raman spectroscopy to determine the phase state of three sulfate micro-inclusions in the Dome Fuji ice core at 2798.5-m depth. Using a temperature-ramp test, we find a peak position at 984 cm<sup>-1</sup> and a change in the full width at half maximum (FWHM) of the S-O stretching mode that identifies the sulfate in the micro-droplets. Considering the peak position and FWHM of sulfate inclusions, we argue that the sulfate would have existed as a micro-droplet liquid on an air hydrate in the ice. The increase in the low frequencies of the Raman spectrum that we detect can be generally used to identify liquids in natural ice. Our investigation also indicates that the surface of air-hydrates in ice may be a preferred location for liquid micro-inclusions.