

## Compound auroral micromorphology: Ground-based high-speed imaging

Ryuho Kataoka<sup>1,2</sup>, Yoko Fukuda<sup>3</sup>, Yoshizumi Miyoshi<sup>4</sup>, Hiroko Miyahara<sup>5</sup>, Satoru Itoya<sup>6</sup>, Yusuke Ebihara<sup>7</sup>, Donald Hampton<sup>8</sup>,  
Hanna Dahlgren<sup>9,10</sup>, Daniel Whiter<sup>11</sup>, and Nickolay Ivchenko<sup>12</sup>

<sup>1</sup>*National Institute of Polar Research, 10-3 Midori-cho, Tachikawa, Tokyo 190-8518, Japan.*

<sup>2</sup>*Department of Polar Science, The Graduate University for Advanced Studies (SOKENDAI)*

<sup>3</sup>*Department of Earth and Planetary Science, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0003, Japan.*

<sup>4</sup>*Solar-Terrestrial Environment Laboratory, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, 464-8601, Japan.*

<sup>5</sup>*Musashino Art University, 1-736 Ogawa-cho, Kodaira-shi, Tokyo 187-8505, Japan.*

<sup>6</sup>*Japan Science Foundation, 2-1 Kitanomaru Park, Chiyoda-ku, Tokyo 102-0091, Japan.*

<sup>7</sup>*Research Institute for Sustainable Humanosphere (RISH), Kyoto University, Gokasho, Uji, Kyoto 611-0011, Japan.*

<sup>8</sup>*Geophysical Institute, University of Alaska Fairbanks, 903 Koyukuk Drive, Fairbanks, AK 99775-7320, USA.*

<sup>9</sup>*School of Electrical Engineering, Royal Institute of Technology KTH, S10044 Stockholm, Sweden.*

<sup>10</sup>*School of Physics and Astronomy, University of Southampton, Southampton, UK.*

<sup>5</sup>*Musashino Art University, 1-736 Ogawa-cho, Kodaira-shi, Tokyo 187-8505, Japan.*

<sup>6</sup>*Finish Meteorological Institute, Helsinki, Finland.*

Auroral microphysics still remains widely unexplored because of the technical limitation of high-speed imaging. Cutting-edge ground-based optical observations using sCMOS cameras recently enabled us to observe the fine-scale morphology of aurora at magnetic zenith for a variety of rapidly varying faint features for long uninterrupted periods. We report two interesting examples of unexpected combinations of fine-scale rapidly varying auroral features as observed by the sCMOS cameras installed at Poker Flat Research Range, Alaska in February 2014. The first example shows that flickering rays and pulsating modulation simultaneously appeared at the middle of a surge in the pre-midnight sector. The second example shows localized flickering aurora associated with growing eddies at the poleward edge of an arc in the midnight sector.