

電離圏へのエネルギー流入と酸素及び水素イオン流出との経験的關係式の太陽天頂角依存性

北村成寿¹、関華奈子¹、桂華邦裕¹、西村幸敏²、堀智昭¹、Robert J. Strangeway³、Eric J. Lund⁴

¹名古屋大学太陽地球環境研究所

²Department of Atmospheric and Oceanic Science, University of California, Los Angeles

³Institute of Geophysics and Planetary Physics, University of California, Los Angeles

⁴Space Science Center, University of New Hampshire

Solar zenith angle dependence of empirical formulas between energy inputs to the ionosphere and O⁺ and H⁺ ion outflows

Naritoshi Kitamura¹, Kanako Seki¹, Kunihiro Keika¹, Yukitoshi Nishimura², Tomoaki Hori¹, Robert J. Strangeway³, and Eric J. Lund⁴

¹Solar-Terrestrial Environment Laboratory, Nagoya University, Nagoya, Japan

²Department of Atmospheric and Oceanic Science, University of California, Los Angeles, CA, USA

³Institute of Geophysics and Planetary Physics, University of California, Los Angeles, CA, USA

⁴Space Science Center, University of New Hampshire, Durham, NH, USA

Recent satellite observations and simulations have clarified that plasma outflows play an important role in abrupt changes in the ion composition in the plasmashet and ring current during geomagnetic storms. Statistical studies by *Strangeway et al.* [2005] and *Brambles et al.* [2011] indicated that fluxes of ion outflows are correlated well with soft electron precipitation (precipitating electron density and electron density in the loss cone), and DC and Alfvénic Poynting fluxes using the data obtained by the FAST satellite near the cusp region in the dayside during the 24-25 September 1998 geomagnetic storm. To evaluate the correlations for H⁺ and O⁺ ions separately, we performed statistical studies using the ion composition data in addition to the ion, electron, and field data obtained by the FAST satellite during January 1998 and January 1999. The longer dataset enables us to identify empirical formulas between outflowing O⁺ and H⁺ ion fluxes and precipitating electron densities, DC and Alfvénic Poynting fluxes in a wide solar zenith angle (SZA) range (45°–145°). These empirical formulas would be useful for global magnetospheric simulations as the boundary conditions. Under dark conditions, H⁺ ion fluxes increase with increasing precipitating electron density, but not as much as those do under sunlit conditions. The precipitating electron density that corresponds to the H⁺ ion flux of $\sim 10^7$ /cm²/s (mapped to 1000 km altitude) decreases with increasing SZA. This SZA dependence is less clear for O⁺ ions as compared with H⁺ ions. The empirical formulas between outflowing O⁺ and H⁺ ion fluxes and DC and Alfvénic Poynting fluxes are not so strongly affected by SZA. Under sunlit conditions, the flux of O⁺ ions tends to be larger than that of H⁺ ions, while H⁺ ions tend to become dominant under dark conditions. Intense ion (especially O⁺ ion) outflow events (number flux larger than $\sim 10^8$ /cm²/s mapped to 1000 km altitude) mostly occurred under sunlit conditions or near the terminator.

References

Brambles, O. J., W. Lotko, B. Zhang, M. Wiltberger, J. Lyon, and R. J. Strangeway, Magnetosphere sawtooth oscillations induced by ionospheric outflow, *Science*, 332, 1183–1186, 2011.

Strangeway, R. J., R. E. Ergun, Y.-J. Su, C. W. Carlson, and R. C. Elphic, Factors controlling ionospheric outflows as observed at intermediate altitudes, *J. Geophys. Res.*, 110, A03221, 2005.