

**Meteorological data from ice-free areas in Yukidori Zawa, Langhovde, Kizahashi Hama,
Skarvsnes and Skallen in Sôya Coast, East Antarctica during 2014–2015**

Sakae KUDOH^{1,2}, Kunio T. TAKAHASHI^{1,2}, Tomoko ISHIHARA³,

Megumu TSUJIMOTO¹, Ryosuke NAKAI⁴, Atsushi C. SUZUKI⁵, Yukiko TANABE^{1,2},

Masaki UCHIDA^{1,2} and Satoshi IMURA^{1,2}

¹ National Institute of Polar Research, Research Organization of Information and Systems,
10–3, Midori-cho, Tachikawa, Tokyo 190-8518.

² Department of Polar Science, School of Multidisciplinary Sciences, SOKENDAI (The Graduate
University for Advanced Studies), 10–3, Midori-cho, Tachikawa, Tokyo 190-8518.

³ Graduate School of Life Science, University of Hyogo, 3–2–1, Koto, Kamigori-cho, Ako-gun,
Hyogo 678-1297.

⁴ National Institute of Genetics, Yata 1111, Mishima, Shizuoka 411-8540.

⁵ Research and Education Center for Natural Science, Keio University, 4–1–1 Hiyoshi, Kohoku-ku,
Yokohama, Kanagawa 223-8521.

*Corresponding author. E-mail: skudoh@nipr.ac.jp

1. Introduction

This report presents meteorological data recorded by automatic weather stations (AWSs) in the Yukidori Zawa (Yukidori Valley), Langhovde, and in Kizahashi Hama, Skarvsnes, and Skallen on the Sôya Coast of East Antarctica, between Japanese Antarctic Research Expedition 55 (JARE-55) in 2014 and JARE-56 in 2015. This study, part of the National Institute of Polar Research (NIPR) project “Monitoring of terrestrial ecosystems (AMB06),” recorded environmental conditions for terrestrial organisms in ice-free areas of Syowa Oasis. The AWSs continuously monitored air temperature, relative humidity, solar radiation, photosynthetically active radiation (PAR), ultraviolet radiation (UV), wind speed and direction, and air pressure, logging them automatically at intervals of 10 min, 1 h, and 1 day.

2. Materials and methods

Observation sites: Yukidori Zawa, in central Langhovde, is Antarctic Specially Protected Area No. 141, where a relatively rich community of vegetation occupies ice-free areas in Syowa Oasis (Kosugi *et al.*, 2014). An AWS was installed at a flat site in the middle of the valley at 69°14'28"S, 39°44'21"E approximately 53 m above sea level, as determined by GPS (Fig. 1). Another AWS was installed at a coastal site in the southeastern part of Kizahashi Hama, a sandy beach in central Skarvsnes, at 69°28'25"S, 39°36'43"E at an elevation of 3 m (Fig. 1). A new AWS was installed at a rocky terrace in the southern coast of Skallen at 69°40'25"S, 39°24'13"E at an elevation of 15 m (Fig. 1).

Observation systems: Each AWS consisted of a data logger (CR1000-4M-XT, Campbell Scientific, USA), an anemometer (05103, Young, USA), a hydrothermometer (HMP155D, Vaissala, Finland), a solar radiation sensor (PCM-01(L), Prede, Japan), a PAR sensor (PAR-01(L), Prede, Japan), a UV sensor (CUV3, Kipp & Zonen, Germany), a barometer (PTB210, Vaissala, Finland) and six sets of lithium ion batteries (TL5930/T, 3.6 V, four cells connected in series for a total of 14 V) capable of supplying 3 years of electric power (19 Ah) to the system. The logger, barometer and batteries were stored in a watertight plastic container (Pelican Products, USA), and the other sensors were mounted on a tripod. Specifications of each sensor had been listed in the previous report (Kudoh *et al.*, 2015). The AWS at Yukidori Zawa was installed on 25 December 2009; that at Kizahashi Hama was installed on 2 February 2010, and that at Skallen was installed on 11 January 2014.

Data logger settings: Every 10 min, the logger compiled a data file containing 10-min average of continuous measurements of wind speed, wind direction, and radiation (solar radiation, PAR and UV), and instantaneous values of temperature, humidity and air pressure. Every 60 min, it compiled 60-min average wind and radiation data with instantaneous values of temperature, humidity and air pressure. In addition, every 24 h it created a summary of daily average wind speed and direction, temperature, humidity, solar radiation, PAR, UV and air pressure. This file also included maximum and minimum values for all parameters and time stamps for all of these values except for solar radiation, PAR and UV. These data were retrieved by the authors during the JARE-56 summer party.

3. Data

We have converted the 10-min, 60-min and 24-h data from the AWS data loggers to CSV files representing each year from 1 January 2014 to 31 January 2015 for Yukidori Zawa, from 1 January 2014 to 10 January 2015 for Kizahashi Hama, and from 12 January 2014 to 25 January 2015 for Skallen, respectively.

We noted during our maintenance visit in January 2014 that the opaque cosine correct plate on the PAR sensor had been broken. The broken sensor had been changed to a new one by the JARE-56 members on 10 January 2015, then, the PAR data during this period had no means. Therefore we removed this portion of the PAR data from Kizahashi Hama in all summary files for 2014 and 2015.

Following is the complete list of data files.

1. Data from Yukidori Zawa AWS

1-1. 10-min summary files

[10min_data_Yukidori_2014.csv](#)

[10min_data_Yukidori_2015.csv](#)

1-2. 60-min summary files

[60min_data_Yukidori_2014.csv](#)

[60min_data_Yukidori_2015.csv](#)

1-3. 24-h summary files

[24hrs_data_Yukidori_2014.csv](#)

[24hrs_data_Yukidori_2015.csv](#)

2. Data from Kizahashi Hama AWS

2-1. 10-min summary files

[10min_data_Kizahashi_2014.csv](#)

[10min_data_Kizahashi_2015.csv](#)

2-2. 60-min summary files

[60min_data_Kizahashi_2014.csv](#)

[60min_data_Kizahashi_2015.csv](#)

2-3. 24-h summary files

[24hrs_data_Kizahashi_2014.csv](#)

[24hrs_data_Kizahashi_2015.csv](#)

3. Data from Skallen AWS

3-1. 10-min summary files

[10min_data_Skallen_2014.csv](#)

[10min_data_Skallen_2015.csv](#)

3-2. 60-min summary files

[60min_data_Skallen_2014.csv](#)

[60min_data_Skallen_2015.csv](#)

3-3. 24-h summary files

[24hrs_data_Skallen_2014.csv](#)

[24hrs_data_Skallen_2015.csv](#)

4. Members who carried out the field study

The AWSs were installed by Sakae Kudoh, Yukiko Tanabe, Masaki Uchida and Takashi Osono (JARE-51 summer), and by Sakae Kudoh, Tomoko Ishihara and Kunio T. Takahashi (JARE-55 summer). Data acquisition was done by Megumu Tsujimoto, Atsushi C. Suzuki and Ryosuke Nakai (JARE-56 summer).

5. Data policy

Before using the data for publication or presentation in any media, please request permission in writing. Inquiries should be addressed to:

Sakae KUDOH

Inter-university Research Institute Corporation Research Organization of Information and Systems,
National Institute of Polar Research, 10-3, Midori-cho, Tachikawa-shi, Tokyo 190-8518, Japan

Telephone +81-42-512-0739 / Facsimile +81-42-528-3492

E-mail skudoh@nipr.ac.jp

Acknowledgements

We thank all the members of JARE-51, JARE-55 and JARE-56, as well as the officers and crew of the icebreaker *Shirase*, for their support.

References

- Kosugi, M., Kurosawa, N., Kawamata, A., Kudoh, S. and Imura, S. (2015); Year-round micrometeorological data from the habitats of terrestrial photosynthetic organisms in Langhovde, East Antarctica, during 2013. JARE data reports, **333** (Terrestrial biology **8**), 99 p.
- Kudoh, S., Tanabe, Y., Uchida, M. and Imura, S. (2015); Meteorological data from ice-free areas in Yukidori Zawa, Langhovde and Kizahashi Hama, Skarvsnes in Sôya Coast, East Antarctica during 2009–2014. JARE data reports, **334** (Terrestrial biology **9**), 7 p.




Fig. 1. Maps of AWS sites. Left panel is a location map of Syowa Oasis (ice-free areas shown in black) on the Sôya Coast, East Antarctica. Right panels indicate the positions of Yukidori (top) and Kizahashi (middle), and Skallen (bottom) AWSs. Locations are outlined in red in the left panel.