

樹木年輪, および生態系モデルを用いた環北極域における 樹木の気候変動に対する応答の時空間変動解析

鄭峻介^{1,2}, 杉本敦子², L. モカン³, 松浦陽次郎⁴, 大沢晃⁵, 米延仁志⁶, 佐藤永⁷, T.マキシモフ⁸
¹ 国立極地研究所, ² 北海道大学, ³ 長江大学, ⁴ 森林総合研究所, ⁵ 京都大学, ⁶ 鳴門教育大学,
⁷ 海洋研究開発機構, ⁸ ロシア寒冷圏生物学研究所

A spatio-temporal pattern of past tree response to climate changes deduced from tree-ring width, delta-¹³C and a DGVM over the pan-Arctic ecosystems

Shunsuke Tei^{1,2}, Atsuko Sugimoto², Liang Maochang³, Yojiro Matsuura⁴, Akira Osawa⁵,
Hitoshi Yonenobu⁶, Hisashi Sato⁷, Trofim Maximov⁸

¹National Institute for Polar research, ²Hokkaido University, ³Yangtze University,
⁴Forestry and Forest Products Research Institute, ⁵Kyoto University, ⁶Naruto University of Education,
⁷JAMSTEC, ⁸Institute for Biological Problems of Cryolithozone SD RSA

Arctic and boreal ecosystems are exposed to rapid and strong increases in temperature and related environmental changes under Arctic amplification. Yet, there is uncertainty how trees in those ecosystems respond to the changes due to an insufficiency of such long term records and this is where tree-rings can provide an advantage. Early dendrochronological studies in the region focused on the positive growth of trees to warmth (D'Arrigo and Jacoby, 1993). However, A number of more recent studies have demonstrated a reduced sensitivity of tree growth to rising temperatures (now referred to as “divergence problem”) at least since the 1960s (e.g., Wilson et al., 2007). Although several studies (e.g., Barber et al., 2000) suggested that temperature-induced drought may limit tree growth under the limited availability of soil moisture, the underlying processes for the phenomenon are not well understood.

We here investigate past tree response to climate changes, especially to warming, using retrospective analyses from tree-ring width and carbon isotope ratios (delta-¹³C) of three genera (*Larix*, *Picea* and *Pinus*) in 6 forest sites with a strong gradient of temperature and precipitation, reaching from northern Europe to northern America; Kalina(59N, 27E), Yakutsk(62N, 129E), Ust'Maya(60N, 133E), Chokurdakh(70N, 148E), Inuvik(68N, 133W) and Fort Smith(60N, 112W). The results suggest that tree response to past climate changes have varied with regions. The tree responses to warming were negative in eastern Siberia forests, resulting in decreasing trend of tree growth over past 60 years. On the other hand, the negative effect of warming is not seen in European and Canadian forests, where no decrease trend of growth is observed. The results then have been used in testing a dynamic global vegetation model (SEIB-DGVM, Sato et al., 2007). The simulated annual net primary productions (NPP) show no decrease trend over the study period and discrepancy from tree-ring based long-term (more than half-decadal) growth variations in eastern Siberian forests, although relatively better reproductions of the model for the variations are obtained in European and Canadian forests. Our results imply that the negative effect of warming override the expected positive effects i.e., warming-induced lengthened growing season and increase in photosynthetic ratio, in arid region such as eastern Siberia, suggesting further reduction of tree growth by future warming. Yet, the negative effect of warming on tree growth (especially for inter-decadal variations) is not well reproduced in the model and therefore the negative effect should be incorporated into future DGVM researches for accurate prediction of future ecosystem functions.