

Odin/OSIRIS データを用いた地磁気活動に対する Na 層応答特性の調査

津田卓雄¹、中村卓司²、江尻省²、西山尚典²、細川敬祐¹、高橋透¹、J. Gumbel³、J. Hedin³

¹ 電気通信大学

² 国立極地研究所

³ ストックホルム大学

Investigation on Na layer response to geomagnetic activities using Odin/OSIRIS data

T. T. Tsuda¹, T. Nakamura², M. K. Ejiri², T. Nishiyama², K. Hosokawa¹, T. Takahashi¹, J. Gumbel³, and J. Hedin³

¹*The University of Electro-Communications*

²*National Institute of Polar Research*

³*Stockholm University*

The Na layer is normally distributed from 80 to 110 km, and the height range is corresponding to the ionospheric D and E region. In the polar region, the energetic particles precipitating from the magnetosphere can often penetrate into the E region and even into the D region. Thus, the influence of the energetic particles to the Na layer is one of interests in the aspect of the atmospheric composition change accompanied with the auroral activity.

There are several previous studies in this issue. For example, recently, we have reported an initial result on a clear relationship between the electron density increase (due to the energetic particles) and the Na density decrease from observational data sets obtained by Na lidar, EISCAT VHF radar, and optical instruments at Tromsø, Norway on 24-25 January 2012. However, all of the previous studies had been carried out based on case studies by ground-based lidar observations.

In this study, we have performed, for the first time, statistical analysis using Na density data from 2004 to 2009 obtained with the Optical Spectrograph and InfraRed Imager System (OSIRIS) onboard Odin satellite. In the presentation, we will show relationship between the Na density and geomagnetic activities, and its latitudinal variation. Based on these results, the Na layer response to the energetic particles will be discussed.