

南極コケ坊主生態系における真核微生物の驚くべき多様性

中井亮佑^{1,2}、長沼毅¹、鹿児島浩³、仁木宏典³、小原雄治³、
伊村智⁴、神田啓史⁴、柳原克彦⁵、馬場知哉⁵、阿部貴志⁶

¹ 広島大学

² 日本学術振興会特別研究員

³ 国立遺伝学研究所

⁴ 国立極地研究所

⁵ 新領域融合研究センター

⁶ 長浜バイオ大学

Unsuspected eukaryotic diversity of aquatic moss pillars inhabiting a freshwater lake in East Antarctica, based on 18S rRNA gene analysis

Ryosuke Nakai^{1,2}, Takeshi Naganuma¹, Hiroshi Kagoshima³, Hironori Niki³, Yuji Kohara³,
Satoshi Imura⁴, Hiroshi Kanda⁴, Katsuhiko Yanagihara⁵, Tomoya Baba⁵, Takashi Abe⁶

¹ Hiroshima University

² JSPS Research Fellow

³ National Institute of Genetics

⁴ National Institute of Polar Research

⁵ Transdisciplinary Research Integration Center

⁶ Nagahama Institute of Bio-Science and Technology

Aquatic mosses of *Bryum* and *Leptobryum* species in association with green algae, diatoms, and cyanobacteria, form unique tower-like pillars of vegetation (“moss pillars”) in Antarctic lakes (Imura et al., 1999). Moss pillars have distinct redox-affected sections, namely aerobic exterior and anaerobic interior. We previously proposed that a “pillar” is structured based on community and habitat of functionally interdependent organisms; hence, a pillar represents a mini-biosphere or bryosphere. Batteries of 16S rRNA genotypes or phylotypes of eubacteria and cyanobacteria, but not archaeobacteria, have been identified in moss pillars (Nakai et al., 2011). Some phylotypes showed pillar-wide distributions, while others were section-specific. This study analyzed near full-length 18S rRNA gene sequences obtained from two whole moss pillars in Lake Hotoke Ike (formerly known as “B-4”), Skarvsnes, in the vicinity of Syowa Station, East Antarctica. Total 28 PCR clone libraries from two whole moss pillars were constructed, and 96 clones from each library (total 2,688 clones) were randomly selected and sequenced. Homology search suggested that most of the sequences were previously unknown but could be assigned to important phyla including Alveolata, Amoebozoa, Chlorophyta, Fungi, Metazoa, Rhizaria, Rhodophyta, and Stramenopiles. We also found unclassified Alveolata, Amoebozoa, Metazoa, and Stramenopiles. In addition, 18S rRNA sequences related to *Leptobryum pyriforme* and unclassified fungi dominated whole moss pillar libraries, with outer aerobic sections further dominated by Ciliophora (genera *Obertruria*, *Onychodromopsis*, and *Oxytricha*) and Tardigrada (genus *Diphascion*) and inner anaerobic sections sub-dominated by unclassified Amoebozoa. The 18S rRNA gene-based profiles demonstrated that redox-related factors may control distribution of physiologically functioning microbial species in a whole moss pillar.

References

- [1] Imura et al. (1999) Benthic moss pillars in Antarctic lakes. *Polar Biology*, 22: 137-140.
- [2] Nakai et al. (in press) Microflorae of aquatic moss pillars in a freshwater lake, East Antarctica, based on fatty acid and 16S rRNA gene analyses. *Polar Biology*, Online First, DOI: 10.1007/s00300-011-1090-2