

# 林冠が疎なアラスカのクロトウヒ林における林床植生の生長期間の推定モデルの開発

永井 信<sup>1</sup>、伊川浩樹<sup>2,3</sup>、中井太郎<sup>3,4</sup>、永野博彦<sup>3</sup>、金 龍元<sup>3</sup>、小林秀樹<sup>1</sup>、Robert C. Busey<sup>3</sup>、鈴木力英<sup>1</sup>

<sup>1</sup> 海洋研究開発機構地球表層物質循環研究分野/北極環境変動総合研究センター

<sup>2</sup> 農業・食品産業技術総合研究機構農業環境変動研究センター

<sup>3</sup> アラスカ大学フェアバンクス校国際北極圏研究センター

<sup>4</sup> 名古屋大学地球水循環研究センター

## Development of phenology model for evaluation of growing season of understory vegetation in an open-canopy black spruce forest in Alaska

Shin Nagai<sup>1</sup>, Hiroki Ikawa<sup>2,3</sup>, Taro Nakai<sup>3,4</sup>, Hirohiko Nagano<sup>3</sup>, Yongwon Kim<sup>3</sup>, Hideki Kobayashi<sup>1</sup>, Robert C. Busey<sup>3</sup> and Rikie Suzuki<sup>1</sup>

<sup>1</sup>*Department of Environmental Geochemical Cycle Research & Institute of Arctic Climate and Environment Research, Japan Agency for Marine-Earth Science and Technology*

<sup>2</sup>*Institute for Agro-Environmental Sciences, National Agriculture and Food Research Organization*

<sup>3</sup>*International Arctic Research Center, University of Alaska Fairbanks*

<sup>4</sup>*Hydrospheric Atmospheric Research Center, Nagoya University*

To deeply understand the interaction between weather and meteorological changes and ecosystem functions (e.g. photosynthesis) in boreal forests in Alaska, we should accurately evaluate the spatio-temporal variability of timings of start (SGS) and end of growing season (EGS) of understory vegetation. Here, (1) we obtained daily canopy surface images in an open-canopy black spruce forest in Alaska (Poker Flat Research Range site; 65°07'24.10"N, 147°29'14.8"W) from spring in 2011 to summer in 2015 and (2) we examined the relationships between daily mean air temperatures and the timings of SGS and EGS of understory vegetation by analyzing RGB values of daily canopy surface images, and made a statistical phenology model based on their relationships. We found that (1) the interannual variability of daily mean air temperatures over 5°C and below 13°C well affected that of the timings of SGS and EGS, respectively, and (2) the estimation accuracy of the timing of SGS was better than that of EGS.