

ATMOSPHERIC NITROGEN DIOXIDE IN ANTARCTICA (ABSTRACT)

Kazuo SHIBASAKI¹, Toshihiro OGAWA² and Naomoto IWAGAMI²

¹*Kokugakuin University, 10-28, Higashi 4-chome, Shibuya-ku, Tokyo 150*

²*Geophysics Research Laboratory, Faculty of Science, University of
Tokyo, 3-1, Hongo 7-chome Bunkyo-ku, Tokyo 113*

Seasonal behavior of stratospheric NO₂ content was determined for the period of March 1983 to January 1984 from ground-based spectroscopic measurements at Syowa Station. The winter minimum of about $1 \times 10^{15} \text{ cm}^{-2}$ and the summer maximum of about $7 \times 10^{15} \text{ cm}^{-2}$ were observed. This seasonal behavior compares well with those observed at northern high latitudes. A rapid increase in column density started at the end of September before the minimum of total ozone in mid-October. This increase in column density seems to correlate with an increase in sunlit time in the stratosphere. Our results imply that both dynamical and photochemical processes may be involved in the Antarctic ozone depletion. The first balloon observation of stratospheric NO₂ profile in Antarctica was made on November 24, 1982 and two observations on November 12 and 20, 1983 at Syowa Station. All observations were concentrated in November, early summer in Antarctica. Two vertical profiles, sunset and sunrise ones, could be deduced in series from one balloon flight. They should be virtually the same rather than corresponding to different profiles for the evening and morning, because, in this season of year, the sunlit condition lasted all day long at the balloon altitude. No quantitative change can be seen in profiles between 1982 and 1983. Above 25 km altitude, three profiles are basically identical to those observed at middle and high latitudes in the northern hemisphere.

(Received February 7, 1987)

DYNAMICAL FACTORS AFFECTING OZONE MIXING RATIOS IN THE ANTARCTIC LOWER STRATOSPHERE (ABSTRACT)

Masato SHIOTANI¹, John C. GILLE² and Lawrence V. LYJAK²

¹*Geophysical Institute, Kyoto University, Kyoto 606*

²*National Center for Atmospheric Research, Boulder, CO 80307, U.S.A.*

This study shows the climatology and interannual variability of dynamical quantities and ozone mixing ratios during the southern hemisphere spring. Analyses are made mainly for six years, from 1979 to 1984, September through December, using the temperature and geopotential height data provided by the National Meteorological Center (NMC) and the ozone mixing ratio data derived from the solar backscatter ultraviolet (SBUV) instrument on board the Nimbus 7.

The zonal mean climatology shows that the coldest temperatures and zonal winds move poleward and downward from September through November, probably in response to wave forcing. A steep decrease in zonal mean ozone mixing ratios is observed around 60°S toward the south pole in September. With time, this high latitude ozone minimum (or "ozone hole") gets shallower in association with minor warmings and a final warming. Climatological synoptic charts in the lower stratosphere show the circumpolar circulation in the geopotential height field and a prominent planetary wave 1 in the temperature and ozone fields. The phases of the temperature and ozone waves in the lower stratosphere are very similar.

The year-to-year variation of the ozone mixing ratio at high latitudes is related to that of the wave activity during the winter and spring. When the wave activity is vigorous, there are weaker

westerlies, higher temperatures and higher ozone mixing ratios at high latitudes. It is possible that the long term change in the lower stratosphere are also related to the long-term trend in wave activity, although the evidence is not clear. Because the wave activity in 1979 was very vigorous, a simple comparison of atmospheric states between the 1979 and other recent years could lead to misleading conclusions on the rate of ozone decrease over the Antarctic. A long-term chemical effect is not precluded.

(Received January 27, 1987)

HETEROGENEOUS REACTIONS RELATED TO ANTARCTIC OZONE HOLE (ABSTRACT)

Shiro HATAKEYAMA¹ and Ming-Taun LEU²

¹*National Institute for Environmental Studies, Yatabe-machi,
Tsukuba-gun, Ibaraki 305*

²*Jet Propulsion Laboratory, California Institute of
Technology, Pasadena, CA 91109, U.S.A.*

Reactions of chlorine nitrate (ClONO₂) with HCl and H₂O have been investigated using a 320-L Pyrex chamber and long-path FT-IR spectroscopy. Both reactions showed highly heterogeneous nature. Obtained upper limit rate constants for homogeneous reactions were 8.4×10^{-21} and 3.4×10^{-21} cm³ molec⁻¹ s⁻¹ for ClONO₂+HCl and ClONO₂+H₂O, respectively, at 296 ± 2 K at 730 torr total pressure. The yield of HNO₃ from both the reactions was 1.05 ± 0.09 and 0.86 ± 0.08 , respectively. Formation of HOCl was confirmed in the latter reaction. No synergistic effect between HCl and H₂O was observed for the reaction with ClONO₂. The kinetic behavior of the reaction ClONO₂+H₂O was well described by simple first-order kinetics while the behavior of the reaction ClONO₂+HCl was found to obey the Langmuir adsorption isotherm.

(Received December 26, 1986)

ATMOSPHERIC CARBON DIOXIDE CONCENTRATION AT SYOWA STATION (69°00'S, 39°35'E), ANTARCTICA (1985) (ABSTRACT)

Haruta MURAYAMA¹, Masayuki TANAKA², Takakiyo NAKAZAWA²,
Sadao KAWAGUCHI³, Takashi YAMANOUCI³, Shuji AOKI³
and Masataka SHIOBARA⁴

¹*Department of Chemistry, Faculty of Education, Yokohama National
University, 156, Tokiwadai, Hodogaya-ku, Yokohama 240*

²*Upper Atmosphere Research Laboratory, Faculty of Science, Tohoku
University, Aramaki Aoba, Sendai 980*

³*National Institute of Polar Research, 9-10, Kaga 1-chome,
Itabashi-ku, Tokyo 173*

⁴*Meteorological Research Institute, 1-1, Nagamine,
Yatabe-machi, Tsukuba-gun, Ibaraki 305*

Since 1984, continuous measurements of the atmospheric CO₂ concentration have been carried out at Syowa Station, Antarctica. Preliminary inspection of the data showed that; (1) a regular