

MARS–EARTH TRANSFER TIME OF LHERZOLITE YAMATO-793605

Otto EUGSTER and Ernst POLNAU

*Physikalisches Institut, University of Bern,
Sidlerstrasse 5, 3012 Bern, Switzerland*

Abstract: We determined the cosmic-ray exposure age of Yamato-793605 based on cosmic-ray produced ^3He , ^{21}Ne , and ^{38}Ar . The average age is 4.4 ± 1.0 Ma and agrees with the exposure ages of the two other martian lherzolites ALH77005 and LEW88516. The similarity in petrology, chemistry, oxygen isotopic composition, and cosmic-ray exposure history shows that the three lherzolitic martian meteorites originate from the same parent magma and were ejected from Mars by the same ejection event. The basaltic shergottites EET79001, QUE94201, Shergotty, and Zagami strongly differ from the three lherzolites in mineralogical and chemical composition and were ejected by events that occurred 1 Ma and 3 Ma (EET79001) later than that responsible for the delivery of the lherzolites. We propose that these differences are sufficiently important to merit distinct meteorite classification and to name ALH77005, LEW88516, and Y-793605 “lherzolites”, in contrast to the “shergottites” EET79001, QUE94201, Shergotty, and Zagami.

1. Introduction

In the framework of a consortium study of martian meteorite Yamato-793605 (hereafter Y-79) organized by H. KOJIMA, P. WARREN, M. MIYAMOTO, and K. YANAI we obtained a sample for the determination of the cosmic-ray exposure age and the abundances of trapped noble gases. Y-79 (recovered mass 16 g) shows strong affinities in petrography and mineral chemistry to two other martian meteorites Allan Hills 77005 (ALH77) and Lewis Cliff 88516 (LEW88) as demonstrated by MIKOUCHI and MIYAMOTO (1996a, b). These meteorites were classified as lherzolitic shergottites, that strongly differ from the basaltic shergottites Elephant Moraine 79001 (EET79), Queen Alexandra Range 94201 (QUE94), Shergotty, and Zagami in mineralogical and chemical composition. Hence these two classes of martian meteorites do not appear to originate from the same parent magma.

Before having analyzed Y-79 we have shown that the above mentioned meteorites originate from three different asteroidal or cometary impact events on Mars (EUGSTER *et al.*, 1997). The time of Mars ejection is given by the sum of the cosmic-ray exposure age and the terrestrial age. The following ejection times were obtained: EET79-0.8 Ma ago, Shergotty, Zagami, and QUE94-2.8 Ma ago, and ALH77 and LEW88-3.8 Ma ago. In this work we shall show that Y-79 yields the same cosmic-ray exposure age (and, assuming a typically brief terrestrial age, also the same ejection time) as the latter two and, thus, belongs to the same ejection event as ALH77 and LEW88. The terrestrial age of Y-79 is presently not known; but because the terrestrial ages of

the martian meteorites are typically relatively short the ejection times are not significantly higher than the cosmic-ray exposure ages.

The basaltic shergottites and, on the other hand, the lherzolitic martian meteorites ALH77, LEW88, and Y-79 do not originate from the same ejection event and, as mentioned above, not from the same parent magma. We shall show in the discussion that the compositional difference between the basaltic shergottites and the lherzolitic martian meteorites is very large, even larger than between the basaltic shergottites and the three nakhlites, another class of martian meteorites. The absence of any genetic relationship between the basaltic shergottites and the lherzolites, except for being martian, does not justify to consider them to be two subgroups of the same class. We, therefore, propose to use the term "lherzolites" for ALH77, LEW88, and Y-79 and not "lherzolitic shergottites".

2. Sample Preparation and Analytical Procedure

We obtained several representative chips with a total weight of 53 mg. The largest chips were gently crushed in a stainless steel mortar and this material was split into two samples of 20.79 mg and 20.53 mg, respectively. Grain sizes were between 0.5 and 3 mm.

The two samples were analyzed using our standard procedure and mass spectrometer system B. For details, such as background and blank corrections see EUGSTER *et al.* (1993). The analytical results are given in Table 1. All errors correspond to a 95% confidence level (2σ errors). Considering the relatively small sample sizes, resulting in possible inhomogeneities and low noble gas amounts, the agreement between the results for the two analyses is satisfactory. In the following discussion we use average values of these two runs.

Table 1. Results of He, Ne, and Ar measurements of Y-793605.

Sample (g)	⁴ He	²⁰ Ne	⁴⁰ Ar	⁴ He	²⁰ Ne	²² Ne	³⁶ Ar	⁴⁰ Ar
	10 ⁻⁸ cm ³ STP/g			³ He	²² Ne	²¹ Ne	³⁸ Ar	³⁶ Ar
0.02079	35.4	1.21	188.6	4.54	0.884	1.211	1.29	557
	±1.2	±0.04	±8.0	±0.05	±0.012	±0.035	±0.10	±40
0.02053	34.9	1.19	197.0	4.72	0.910	1.241	1.15	593
	±1.1	±0.04	±10.0	±0.09	±0.020	±0.080	±0.10	±55
Average	35.2	1.20	192.8	4.63	0.897	1.226	1.22	575
	±1.0	±0.04	±5.0	±0.10	±0.015	±0.015	±0.07	±20

3. Cosmic-ray Produced Noble Gases

Helium. If ${}^4\text{He}/{}^3\text{He}=5$ (HEYMANN, 1967) is adopted for cosmogenic He, the low ${}^4\text{He}/{}^3\text{He}$ ratio of 4.54 and 4.72 (Table 1) excludes both trapped He and radiogenic ${}^4\text{He}$ contributions. Thus, we conclude that all ${}^3\text{He}$ and ${}^4\text{He}$ are of cosmogenic origin.

Neon. Adopting $({}^{20}\text{Ne}/{}^{21}\text{Ne})_c=0.9$ and non-cosmogenic Ne to have terrestrial atmospheric isotopic composition we obtain ${}^{21}\text{Ne}_c$, ${}^{20}\text{Ne}_{tr}$ and $({}^{22}\text{Ne}/{}^{21}\text{Ne})_c$ as given in




Fig. 1. Ejection times (sum of cosmic-ray exposure age and terrestrial age) for martian meteorites dated until now. For references see EUGSTER *et al.* (1997).

Thus, the ejection of Y-79 from Mars occurred 4–5 Ma ago. This time agrees within errors with that observed for ALH77 and LEW88. The three lherzolites yield an average ejection age of about 4.0 ± 0.5 Ma whereas that for the shergottites QUE94201 (2.91 Ma), Shergotty (2.71 Ma) and Zagami (2.76 Ma) is 2.79 ± 0.12 Ma (EUGSTER *et al.*, 1997). Figure 1 shows the ejection times for all martian meteorite classes, demonstrating that at least five ejection events are responsible for the production of the recovered martian meteorites. It is presently not clear whether the nakhlites and Chassigny represent one or two ejection events.

5. The Lherzolites Should Not Be Called “Shergottites”

The martian meteorite classes differ in mineral and, consequently, in chemical composition (*cf.*, MCSWEEN, 1994). The basaltic shergottites EET79, QUE94, Shergotty and Zagami consist predominantly of the clinopyroxenes pigeonite and augite, whereas the lherzolic martian meteorites ALH77, LEW88 and Y-79 mainly consist of olivine, chromite, and orthopyroxene. In order to quantify the differences in composition (in wt%) between the martian meteorite classes we plot in Fig. 2 the Mg/Ti *versus* the Mg/Al and the Mg/Ca *versus* the Mg/Al ratios. The rectangles shown for the shergottites, lherzolites, and nakhlites represent the range of all values for the members of the respective class. Obviously, the three lherzolic meteorites ALH77, LEW88, and Y-79 are distinct from the basaltic shergottites by factors of about seven for the three ratios shown in Fig. 2. The difference in chemical composition between the lherzolites and the basaltic shergottites is even larger than that between the nakhlites and the basaltic shergottites. Furthermore, on the oxygen isotope fractionation line for martian meteorites all three lherzolites plot in a range for $\delta^{18}\text{O} = 3.88\text{--}4.14$ and $\delta^{17}\text{O} = 2.18\text{--}2.39$, that is, below the range for the basaltic shergottites with $\delta^{18}\text{O} = 4.23\text{--}4.97$ and $\delta^{17}\text{O} = 2.31\text{--}2.76$ (CLAYTON and MAYEDA, 1996). Finally, we conclude from the noble gas isotopic studies that the Mars ejection times of the basaltic shergottites and




Fig. 2. (a) Mg/Ti versus Mg/Al , (b) Mg/Ca versus Mg/Al for martian meteorites. The rectangles indicate the range of the elemental ratios for the meteorites of each class. Data from TREIMAN *et al.* (1986), DREIBUS *et al.* (1992), MITTFELDLT (1994), and WARREN *et al.* (1996).

the lherzolites do not overlap: the former ones were ejected in two events—0.8 Ma (EET79) and 2.8 ± 0.1 Ma (QUE94, Shergotty, and Zagami), whereas the latter ones (ALH77, LEW88, and Y-79) in a earlier event 4.0 ± 0.5 Ma ago. Hence, we propose

to name the two classes “shergottites” and “lherzolites”, respectively.

6. Is There a ^{40}Ar Excess in Y-793605?

Clear evidences for entrapped martian atmospheric gases characterized by an excess of ^{40}Ar and $^{40}\text{Ar}/^{36}\text{Ar}$ ratios >1600 were observed in EET79 (BOGARD and JOHNSON, 1983; BECKER and PEPIN, 1984; SWINDLE *et al.*, 1986) and in Zagami (MARTI *et al.*, 1995). If the K concentration of Y-79 is in the range of those for the other lherzolites (Table 4) a small ^{40}Ar excess is present, even assuming a gas retention age of 1300 Ma (igneous formation age of lherzolite ALH77, SHIH *et al.*, 1982). If the radiogenic ^{40}Ar is $15 \times 10^{-8} \text{ cm}^3 \text{ STP/g}$ (180 Ma gas retention age, JONES, 1986 and McSWEEN, 1994) the $(^{40}\text{Ar}/^{36}\text{Ar})_{\text{r}}$ would be 1000, indicating martian atmospheric Ar. On the other hand, if $^{40}\text{Ar}_{\text{r}} = 155 \times 10^{-8} \text{ cm}^3 \text{ STP/g}$ (1300 Ma gas retention age) the resulting $(^{40}\text{Ar}/^{36}\text{Ar})_{\text{r}} = 213$, close to terrestrial atmospheric $^{40}\text{Ar}/^{36}\text{Ar} = 296$.

Table 4. Excess ^{40}Ar in lherzolites.

Meteorite	K ppm	$^{40}\text{Ar}_{\text{meas}}$	$^{40}\text{Ar}_{\text{r}}$ produced in		Excess ^{40}Ar to	
			180 Ma	1300 Ma	180 Ma	1300 Ma
			$10^{-8} \text{ cm}^3 \text{ STP/g}$			
Y-793605	210	193	15	155	178	38
ALH77005	224	137	16	166	121	0
LEW88516	199	781	15	147	766	634

Acknowledgments

We thank the National Institute of Polar Research for the sample of martian meteorite Yamato-793605 and H. KOJIMA, P. WARREN, M. MIYAMOTO, and K. YANAI for organizing the consortium study. We acknowledge the careful and helpful reviews by Y. MIURA and K. NAGAO. M. ZUBER assisted in the sample preparation, P. GUGGISBERG in mass spectrometry, and B. BALSIGER in the preparation of the manuscript. This work was supported by the Swiss National Science Foundation.

References

- BECKER, R. H. and PEPIN, R. O. (1984): The case for a martian origin of the shergottites: Nitrogen and noble gases in EETA 79001. *Earth Planet. Sci. Lett.*, **69**, 225–242.
- BECKER, R. H. and PEPIN, R. O. (1993): Nitrogen and noble gases in a glass sample from the LEW88516 shergottite. *Meteoritics*, **28**, 637–640.
- BOGARD, D. D. and GARRISON, D. H. (1993): Noble gases in LEW88516 shergottite: Evidence for exposure age pairing with ALH77005. *Lunar and Planetary Science XXIV*. Houston, Lunar Planet. Inst., 139–140.
- BOGARD, D. D. and JOHNSON, P. (1983): Martian gases in an antarctic meteorite? *Science*, **221**, 651–654.
- BOGARD, D. D., NYQUIST, L. E. and JOHNSON, P. (1984): Noble gas contents of shergottites and implications for the Martian origin of SNC meteorites. *Geochim. Cosmochim. Acta*, **48**, 1723–1739.
- CLAYTON, R. N. and MAYEDA, T. K. (1996): Oxygen isotope studies of achondrites. *Geochim. Cosmochim. Acta*, **60**, 1999–2017.

- DREIBUS, G., JOCHUM, K. H., PALME, H., SPETTEL, B., WLOTZKA, F. and WÄNKE, H. (1992): LEW88516: A meteorite compositionally close to the "Martian mantle". *Meteoritics*, **27**, 216.
- EUGSTER, O. and MICHEL, Th. (1995): Common asteroid break-up events of eucrites, diogenites, and howardites and cosmic-ray production rates for noble gases in achondrites. *Geochim. Cosmochim. Acta*, **59**, 177–199.
- EUGSTER, O. and WEIGEL, A. (1992): Exposure histories of lodranites, shergottite LEW88516 and CK-chondrites. *Meteoritics*, **27**, 219.
- EUGSTER, O., MICHEL, Th., NIEDERMANN, S., WANG, D. and YI, W. (1993): The record of cosmogenic, radiogenic, fissionogenic, and trapped noble gases in recently recovered Chinese and other chondrites. *Geochim. Cosmochim. Acta*, **57**, 1115–1142.
- EUGSTER, O., WEIGEL, A. and POLNAU, E. (1997): Ejection times of martian meteorites. *Geochim. Cosmochim. Acta*, **61**, 2749–2757.
- HEYMANN, D. (1967): On the origin of hypersthene chondrites: Ages and shock effects of black chondrites. *Icarus*, **6**, 189–221.
- JONES, J. H. (1986): A discussion of isotopic systematics and mineral zoning in the shergottites: Evidence for a 180 m.y. igneous crystallization age. *Geochim. Cosmochim. Acta*, **50**, 969–977.
- MARTI, K., KIM, J. S., THAKUR, A. N., MCCOY, T. J. and KEIL, K. (1995): Signatures of the martian atmosphere in glass of the Zagami meteorite. *Science*, **267**, 1981–1984.
- McSWEEN, H. J., Jr. (1994): What we have learned about Mars from SNC meteorites. *Meteoritics*, **29**, 757–779.
- MIKOUCHI, T. and MIYAMOTO, M. (1996a): A new member of lherzolic shergottite from Japanese Antarctic Meteorite Collection: Mineralogy and petrology of Yamato-793605. *Antarctic Meteorites XXI*. Tokyo, Natl Inst. Polar Res., 104–106.
- MIKOUCHI, T. and MIYAMOTO, M. (1996b): Comparative mineralogy of antarctic lherzolic shergottites ALH77005, LEW88516, and Y793605. *Meteorit. Planet. Sci.*, **31**, Suppl., A89–A90.
- MITTFELHLDT, D. W. (1994): ALH84001, a cumulate orthopyroxenite member of the martian meteorite clan. *Meteoritics*, **29**, 214–221.
- NISHIZUMI, K. and CAFFEE, W. (1996): Exposure history of shergottite Queen Alexandra Range 94201. *Lunar and Planetary Science XXVII*. Houston, Lunar Planet. Inst., 961–962.
- OTT, U. and LÖHR, H. P. (1992): Noble gases in the new shergottite LEW88516. *Meteoritics*, **27**, 271.
- SHIH, C.-Y., NYQUIST, L. E., BOGARD, D. D., MCKAY, G. A., WOODEN, J. L., BANSAL, B. M. and WIESMANN, H. (1982): Chronology and petrogenesis of young achondrites, Shergotty, Zagami, and ALHA77005: Late magmatism on a geologically active planet. *Geochim. Cosmochim. Acta*, **46**, 2323–2344.
- SWINDLE, T. D., CAFFEE, M. W. and HOHENBERG, C. M. (1986): Xenon and other noble gases in shergottites. *Geochim. Cosmochim. Acta*, **50**, 1001–1015.
- TREIMAN, A. H., DRAKE, M. J., JANSSENS, M.-J., WOLF, R. and EBHARA, M. (1986): Core formation in the Earth and Shergottite Parent Body (SPB): Chemical evidence from basalts. *Geochim. Cosmochim. Acta*, **50**, 1071–1091.
- WARREN, P., KALLEMEYN, G. W., ARAI, T. and KANEDA, K. (1996): Compositional-petrologic investigation of eucrites and the QUE94201 shergottite. *Antarctic Meteorites XXI*. Tokyo, Natl Inst. Polar Res., 195–197

(Received February 10, 1997; Revised manuscript accepted April 24, 1997)