

—研究論文—  
Scientific Paper

## Evaluation of SO<sub>2</sub> Emission from the 1982 Eruption of El Chichon by Glaciological and Satellite Methods

Mika KOHNO<sup>1\*</sup>, Minoru KUSAKABE<sup>1</sup> and Yoshiyuki FUJII<sup>2</sup>

南極の積雪試料を用いたエル・チチョン火山 1982 年噴火に伴う  
二酸化硫黄放出量の推定

河野美香<sup>1\*</sup>・日下部 実<sup>1</sup>・藤井理行<sup>2</sup>

**要旨:** 火山噴火に伴う大気中への硫黄放出量の推定方法の一つに、極地方の氷床の硫酸堆積量に基づく方法がある。これは、氷床に堆積した雪に含まれる火山性硫酸の量にスケールリングファクター（核実験により放出され成層圏を経て降下した放射性物質の広がり方に基づいて決められた、成層圏を経由する物質運搬効率を地球上の任意の地点で与えるファクター）を乗じる方法である。ケーススタディとして、近年の爆発的噴火の一つであるメキシコのエルチチョン火山 1982 年噴火について、南極氷床のドームふじ雪試料から硫黄放出量を推定した。同噴火による放出量は、人工衛星に搭載した紫外線分光器（TOMS）による噴煙のリモートセンシングにより推定されており（7 Mt SO<sub>2</sub>）、本研究による推定値はこれの 2.5 倍程度の値であった（17 Mt SO<sub>2</sub>）。このことは、過去の大噴火に伴う硫黄放出量の推定に、氷床の硫酸堆積量に基づく方法が適用できることを示している。

**Abstract:** The SO<sub>2</sub> emission at the time of the 1982 eruption of El Chichon in Mexico was estimated by a glaciological method based on sulfate ion concentration and bomb-produced total  $\beta$  activity in snow collected in Antarctica. The glaciological estimate of the SO<sub>2</sub> emission is 17 ± 1 Mt SO<sub>2</sub>, somewhat greater than the spectroscopic estimate (7 Mt SO<sub>2</sub>) measured by a satellite-carried total ozone mapping system. We conclude that the glaciological method is useful to estimate SO<sub>2</sub> emissions of past explosive volcanic eruptions in the equatorial region.

### 1. Introduction

Major explosive volcanic eruptions emit a large amount of sulfur dioxide (SO<sub>2</sub>) into the stratosphere. The SO<sub>2</sub> is gradually converted into sulfuric acid aerosols in the stratosphere, spreads over the globe, and eventually falls onto the Earth's surface in a few years. Snow that accumulated continuously on polar ice sheets contains a record of such volcanic sulfate signals. The stratospheric sulfuric acid aerosols reduce atmospheric transmissivity and temperature at the Earth's surface. The effects on climate of

<sup>1</sup> 岡山大学固体地球研究センター。Institute for Study of the Earth's Interior, Okayama University, Misasa, Tottori 682-0194.

<sup>2</sup> \* (Present address) 国立極地研究所。National Institute of Polar Research, Kaga 1-chome, Itabashi-ku, Tokyo 173-8515.

recent explosive eruptions such as Agung in 1963, St. Helens in 1980, El Chichon in 1982, and Pinatubo in 1991 have been well-documented (DEVINE *et al.*, 1984; RAMPINO and SELF, 1984; MINNIS *et al.*, 1993; SELF *et al.*, 1996). For evaluation of the relationship between volcanism in the past and subsequent climatic change, it is essential to estimate the amount of SO<sub>2</sub> emitted by past explosive volcanic eruptions.

Volcanic SO<sub>2</sub> emission has been estimated by (1) the glaciological method, (2) the petrological method, and (3) the spectroscopic method. The glaciological method is based on the SO<sub>4</sub><sup>2-</sup> ion concentration in a polar snow sample that accumulated at the time of a volcanic eruption, multiplied by the local accumulation rate of snow and the ratio of locally measured bomb-produced total  $\beta$  activity to the total  $\beta$  activity produced at the time of atmospheric bomb tests in the equatorial region in 1952 and 1954 (HAMMER *et al.*, 1980; CLAUSEN and HAMMER, 1988; LANGWAY *et al.*, 1988; ZIELINSKI, 1995). The petrological method is based on the difference of sulfur concentrations between glass inclusions in phenocrysts and matrix glasses scaled by the mass of the magma erupted. Glass inclusions are considered to preserve pre-eruptive dissolved volatile concentration in melt of the magma, whereas matrix glasses have lost the dissolved volatiles during decompression and eruption of the magma (SIGURDSSON, 1982; DEVINE *et al.*, 1984; PALAIS and SIGURDSSON, 1989; KOHNO, 1992; MANDEVILLE *et al.*, 1996). The spectroscopic method, which has been established as a direct method to measure SO<sub>2</sub> emission, is based on UV spectrometric measurement in the volcanic cloud by a satellite-carried total ozone mapping system (TOMS: KRUEGER, 1983; KRUEGER *et al.*, 1990; BLUTH *et al.*, 1992).

Recently it has been pointed out, particularly since spectroscopic measurements have become available, that the petrological estimates are generally much lower than the spectroscopic estimates by a factor of 10 or more, the spectroscopic method being regarded as providing plausible estimates of SO<sub>2</sub> emission (ANDRES *et al.*, 1991; WESTRICH and GERLACH, 1992; GERLACH *et al.*, 1994; KAZAHAYA and SHINOHARA, 1994). The difference is referred to as excess sulfur degassing. The cause of excess sulfur degassing is currently one of the most active research topics in volcanology (DEVINE *et al.*, 1984; RAMPINO and SELF, 1984; KRUEGER *et al.*, 1990; ANDRES *et al.*, 1991; WALLACE and GERLACH, 1994; GERLACH *et al.*, 1994; HATTORI, 1996; RUTHERFORD and DEVINE, 1996; KRESS, 1997). Because of their own nature, the glaciological and petrologic methods are only used to estimate SO<sub>2</sub> emission by historic volcanic eruptions.

In this study, we compare the spectroscopic and glaciological estimates of volcanic SO<sub>2</sub> emission by the 1982 eruption of El Chichon in Mexico. The eruption was one of the most explosive in recent years. The amount of SO<sub>2</sub> emission estimated by the spectroscopic and petrologic methods have been reported to be 7 Mt SO<sub>2</sub> (BLUTH *et al.*, 1993) and 0.07 Mt SO<sub>2</sub> (DEVINE *et al.*, 1984), respectively, although the amount of the emission estimated by the glaciological method has not been reported yet. We estimated the SO<sub>2</sub> emission by the glaciological method using snow samples collected at Dome Fuji Station, Antarctica.

## 2. Samples and Analytical Procedures

Snow samples were collected in January 1997 from the wall of a 3.8 m deep pit

excavated at Dome Fuji Station (77° 19' 01" S, 39° 42' 12" E, 3810 m a.s.l.) located at the summit of the inland plateau in Queen Maud Land, East Antarctica (Fig. 1). Characteristic of snow stratification observed on the pit-wall was uniform bedding with well-developed depth hoar. Snow samples were collected from 57 layers between surface and 3.75 m depth and stored in 100 ml polyethylene bottles. The stratigraphy of the upper 36 snow layers is shown in Fig. 2a.

The annual average accumulation rate at Dome Fuji Station has been reported to be  $32 \text{ kg} \cdot \text{m}^{-2} \cdot \text{a}^{-1}$  (KAMIYAMA *et al.*, 1989). This was estimated from the tritium peak of 1964/65 which appeared at 0.63 m in water equivalent depth of snow collected at DF-80, 2 km southwest from Dome Fuji Station, in December 1985 (KAMIYAMA *et al.*, 1989). Snow layers of the pit collected in January 1997 were dated using the annual accumulation rate as shown in Fig. 2.

The snow samples, preserved in a low-temperature room ( $-20^\circ \text{C}$ ), were melted in a refrigerator ( $+5^\circ \text{C}$ ) overnight. Concentrations of anions ( $\text{Cl}^-$ ,  $\text{F}^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{NO}_2^-$  and  $\text{NO}_3^-$ ) and cations ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{NH}_4^+$ ,  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$ ) were simultaneously measured using an ion chromatograph (Dionex model DX-500) equipped with AG11 and CG14 guard columns, AS11 and CS14 separator columns, and ASRS-1 and CSRS-1 auto suppressors to concentrate the ionic species having a concentration level of  $\sim 10 \text{ ng} \cdot \text{g}^{-1}$ . The precision and detection limit of the measurement for each ion are




Fig. 1. Location of Dome Fuji Station in Queen Maud Land, East Antarctica.




Fig. 2. (a) Stratigraphy and (b) non-sea-salt sulfate ( $\text{nssSO}_4^{2-}$ ) concentration of the upper 36 snow layers of a 3.8 m deep snow pit excavated in January 1997 at Dome Fuji Station, East Antarctica. In the stratigraphic diagram, solid and dotted lines indicate layer boundaries with and without ice crust, respectively. Hatched shade indicates the degree of hardness (a harder layer is darker).

described in WATANABE *et al.* (1997).

### 3. Basic Principle of the Glaciological Method

The total amount of sulfur from a source volcano can be estimated under the following assumption. The ratio of the amount of sulfur reaching the snow pit site to sulfur emitted from a source volcano in the equatorial region is assumed to be equal to the ratio of total  $\beta$  activity observed at the pit site to the total  $\beta$  activity produced by atmospheric bomb tests in the same region (HAMMER, 1977; HAMMER *et al.*, 1980). This is expressed by the following equation:

$$\text{SO}_2 = [\text{SO}_4]_{\text{snow}}^* \cdot \frac{\beta^{\text{test region}}}{\beta^{\text{pit site}}} \cdot \frac{W_{\text{SO}_2}}{W_{\text{H}_2\text{SO}_4}}, \quad (1)$$

where  $[\text{SO}_4]_{\text{snow}}^*$  is the amount of volcanic sulfate (*i.e.* non-sea-salt sulfate as described later) deposition in snow sample,  $\beta^{\text{test region}}$  is the amount of total  $\beta$  activity produced by the atmospheric nuclear bomb tests and  $\beta^{\text{pit site}}$  is the amount of total  $\beta$  activity measured

at the pit site.  $W_{\text{SO}_2}$  and  $W_{\text{H}_2\text{SO}_4}$  denote the molecular weights of SO<sub>2</sub> and H<sub>2</sub>SO<sub>4</sub>, respectively.

The total  $\beta$  activity consists mainly of <sup>90</sup>Sr and <sup>137</sup>Cs (CLAUSEN and HAMMER, 1988). The basic concept behind this assumption is that the transport of aerosols is controlled by the global atmospheric circulation and that the behavior of sulfuric acid aerosols of volcanic origin and the bomb-produced aerosols are the same on a global scale.

The total  $\beta$  activity at the pit site also needs to be measured. According to the UNSCEAR Report (1982), the total  $\beta$  activity from the US atmospheric bomb tests at 11° N in 1952 and 1954 was 9.56 MCi. Then, sulfur (in terms of SO<sub>2</sub>) emission by volcanic eruption in the equatorial region can be estimated by eq. (1).

The total fallout of sulfuric acid (H<sub>2</sub>SO<sub>4</sub>) in a unit area at a pit site is obtained by multiplying the H<sub>2</sub>SO<sub>4</sub> concentration of a layer, which includes non-sea-salt sulfate ion (nssSO<sub>4</sub><sup>2-</sup>) originating from a volcanic eruption, by the water equivalent thickness of the layer. The nssSO<sub>4</sub><sup>2-</sup> concentration is calculated using the following equation, assuming that Na<sup>+</sup> ions are entirely derived from sea spray:

$$(\text{SO}_4^{2-})_{\text{non-sea-salt}} = (\text{SO}_4^{2-}) - (\text{SO}_4^{2-}/\text{Na}^+)_{\text{seawater}} \cdot (\text{Na}^+), \quad (2)$$

where (SO<sub>4</sub><sup>2-</sup>) and (Na<sup>+</sup>) are the measured SO<sub>4</sub><sup>2-</sup> and Na<sup>+</sup> concentrations of the snow sample, and (SO<sub>4</sub><sup>2-</sup>/Na<sup>+</sup>)<sub>seawater</sub> is the ratio of SO<sub>4</sub><sup>2-</sup> to Na<sup>+</sup> concentrations of seawater being 0.25 in a weight ratio.

#### 4. Estimate of SO<sub>2</sub> Emission from the 1982 Eruption of El Chichon Volcano

We show the concentrations of Na<sup>+</sup>, SO<sub>4</sub><sup>2-</sup> and nssSO<sub>4</sub><sup>2-</sup> in Table 1. Two significant nssSO<sub>4</sub><sup>2-</sup> signals were found in layers between 0.06 and 0.12 m depths and between 0.43 and 0.45 m depths, which correspond to 1993 and 1983, respectively (Fig. 2b). Judging from its date, the deeper signal is likely due to the deposition of sulfuric acid aerosols from the 1982 eruption of El Chichon volcano (in Mexico, 17° 36' N, 93° 23' W). In 1982, El Chichon volcano emitted 7 Mt of SO<sub>2</sub> to the stratosphere, high enough to give a clear nssSO<sub>4</sub><sup>2-</sup> signal in polar regions (BLUTH *et al.*, 1993; KOHNO *et al.*, 1996). The nssSO<sub>4</sub><sup>2-</sup> signal found between 0.06 and 0.12 m depths may include sulfuric acid aerosols originating from the 1991 eruption of Pinatubo volcano (in the Philippines, 15° 14' N, 120° 35' E) and the 1991 eruption of Cerro Hudson (in Chile, 46° 17' S, 72° 92' W). BLUTH *et al.* (1993) reported the SO<sub>2</sub> emission measured by TOMS during the eruption of Pinatubo volcano to be 20 Mt SO<sub>2</sub> and that of Cerro Hudson volcano to be 2 Mt SO<sub>2</sub>. The Cerro Hudson volcano is closer to Antarctica and thereby yielded substantial amounts of nssSO<sub>4</sub><sup>2-</sup> at the pit site. Since the signal cannot be easily decomposed into the Pinatubo and Cerro Hudson components, we discuss only the emission from the 1982 eruption of El Chichon here.

The deposition of nssSO<sub>4</sub><sup>2-</sup> in the layer between 0.43 and 0.45 m depths, corresponding to 1983, is 4.2 kg·km<sup>-2</sup>. The deposition of nssSO<sub>4</sub><sup>2-</sup>, due to the eruption of El Chichon, was obtained as 2.1 kg·km<sup>-2</sup> by subtracting the background nssSO<sub>4</sub><sup>2-</sup> (2.1 kg·km<sup>-2</sup>) originating from oxidation of dimethylsulfide (DMS) produced by ocean biogenic activity. The background nssSO<sub>4</sub><sup>2-</sup> level was obtained by averaging all layers

Table 1. Chemical composition of snow from the surface to 1.9 m depth in a snow pit at Dome Fuji Station.

Water equivalent in m		Na <sup>+</sup>	SO <sub>4</sub> <sup>2-</sup>	nssSO <sub>4</sub> <sup>2-</sup>
Upper depth	Lower depth	ng · g <sup>-1</sup>	ng · g <sup>-1</sup>	ng · g <sup>-1</sup>
0.00	0.01	1.8	32.8	32.3
0.01	0.02	4.6	61.1	59.9
0.02	0.04	11.4	85.9	83.1
0.04	0.05	8.3	86.7	84.6
0.05	0.06	142.2	127.6	124.0
0.06	0.08	21.9	204.1	198.6
0.08	0.11	21.3	231.1	225.7
0.11	0.12	43.3	247.5	236.6
0.12	0.13	3.8	161.7	160.8
0.13	0.14	2.1	108.0	107.5
0.14	0.15	32.2	125.9	117.8
0.15	0.17	27.5	123.8	116.9
0.17	0.19	14.2	96.6	93.1
0.19	0.22	42.5	141.4	130.8
0.22	0.23	3.2	79.0	78.1
0.23	0.26	15.0	124.7	121.0
0.26	0.27	32.7	112.7	104.5
0.27	0.29	1.1	41.5	41.3
0.29	0.30	1.9	41.5	41.0
0.30	0.31	8.9	95.9	93.7
0.31	0.33	16.6	109.7	105.5
0.33	0.34	15.1	108.0	104.2
0.34	0.36	15.0	109.1	105.3
0.36	0.37	40.5	172.0	161.9
0.37	0.39	23.7	117.7	111.8
0.39	0.42	15.9	98.2	94.2
0.42	0.43	30.1	113.3	105.8
0.43	0.45	32.4	116.7	108.6
0.45	0.48	11.4	217.1	214.2
0.48	0.49	20.8	139.1	133.9
0.49	0.50	1.8	63.4	63.0
0.50	0.51	18.5	80.7	76.1
0.51	0.53	19.3	102.1	97.3
0.53	0.55	33.6	97.3	88.9
0.55	0.57	26.9	88.5	81.8
0.57	0.59	24.5	105.4	99.2

except the volcanic layers (Table 2 and Fig. 2b).

Deposition of the total  $\beta$  activity, derived from nuclear tests in 1952 and 1954, at DF-80 near Dome Fuji Station was found to be  $2.84 \times 10^{-1} \text{ mCi} \cdot \text{km}^{-2}$  by integrating the total  $\beta$  activity in the layers between 0.83 and 1.04 m in the snow pit in 1985. These




Fig. 3. Vertical profile of total  $\beta$  activity of snow pit in December 1985 at DF-80 near Dome Fuji Station (KAMIYAMA *et al.*, 1997). Unit of dpm means disintegration per minute ( $1 \text{ dpm} = 1/3.7 \times 10^{-7} \text{ mCi}$ ). A hatched area indicates a layer of total  $\beta$  activity produced by the nuclear tests in 1952 and 1954.

depths correspond to the middle of 1953 and the end of 1959, respectively (Fig. 3; data after KAMIYAMA *et al.*, 1997). These dates were determined by considering a time difference of  $\sim 3$  years (POURCHET and PINGLOT, 1979) between nuclear tests and appearance of the total  $\beta$  activity peak on Antarctica, and their residence time in the stratosphere of  $\sim 1.5$  years (POURCHET *et al.*, 1983). The  $\beta^{\text{pit site}}$  value in eq. (1) was corrected for radioactive decay between the time of deposition and measurement under the assumption that the total  $\beta$  activity was due only to  $^{90}\text{Sr}$  ( $T_{1/2} = 28$  years) and  $^{137}\text{Cs}$  ( $T_{1/2} = 29$  years). The corrected  $\beta^{\text{pit site}}$  is calculated to be  $7.53 \times 10^{-1} \text{ mCi} \cdot \text{km}^{-2}$  assuming that the total  $\beta$  activity was mainly due to  $^{90}\text{Sr}$ . The  $\beta^{\text{test region}}$  has been reported to be  $9.56 \times 10^9 \text{ mCi}$  as mentioned above (UNSCEAR Report, 1982; CLAUSEN and HAMMER, 1988). Thus, a scaling factor, the ratio of  $\beta^{\text{test region}}$  to  $\beta^{\text{pit site}}$ , is calculated to be  $1.3 \times 10^{10} \text{ km}^2$  for the region around Dome Fuji Station (Table 2).

The amount of SO<sub>2</sub> emission from the 1982 eruption of El Chichon is thus

Table 2. Glaciological SO<sub>2</sub> estimate of the 1982 eruption of El Chichon.

Total $\text{kg} \cdot \text{km}^{-2}$	SO <sub>4</sub> deposition Background $\text{kg} \cdot \text{km}^{-2}$	Volcanic $\text{kg} \cdot \text{km}^{-2}$	$\beta^{\text{test region}} / \beta^{\text{pit site}}$ $\text{km}^2$	SO <sub>2</sub> <sup>GE</sup> Mt	SO <sub>2</sub> <sup>SE</sup> Mt
4.2	2.1	2.1	$1.3 \times 10^{10}$	17	7

GE: glaciological estimate; SE: spectroscopic estimate (BLUTH *et al.*, 1993).

Note:  $\beta^{\text{test region}} = 9.56 \times 10^9 \text{ mCi}$  (CLAUSEN and HAMMER, 1988);  $\beta^{\text{pit site}} = 7.53 \times 10^{-1} \text{ mCi}$  after correction for radioactive decay.

calculated to be  $17 \pm 1$  Mt  $\text{SO}_2$  using eq. (1). The error is propagated from the following factors. The error in layer thickness is considered to be within 0.5%. Analytical error of the  $\text{SO}_4^{2-}$  determination with ion chromatography has been reported to be within 0.3% (IGARASHI *et al.*, 1998). The error for the total  $\beta$  activity determination is estimated to be within 7% for the activity level of  $3 \text{ dpm} \cdot \text{km}^{-2}$  (KAMIYAMA, pers. commun.), the mean total  $\beta$  activity emitted in 1952 and 1954. The error in the estimation of the amount of total  $\beta$  activity produced by bomb-tests has not been reported.

The glaciological estimate of the  $\text{SO}_2$  emission of  $17 \pm 1$  Mt  $\text{SO}_2$  is apparently greater than the spectroscopic estimate of 7 Mt  $\text{SO}_2$  (BLUTH *et al.*, 1993). According to SYMONDS *et al.* (1994), the spectroscopic estimate which was directly measured in the volcanic cloud by a satellite-carried TOMS is considered to be an underestimate because low  $\text{SO}_2$  concentration at the margin of the volcanic cloud is below the detection limit of TOMS measurement.

While much work remains to be done to determine the reason for the difference between glaciological and spectroscopic estimates (17 and 7 Mt  $\text{SO}_2$ , respectively), the ratio between glaciological and spectroscopic estimates ( $\sim 2.5$ ) is much less than the ratio ( $\sim 100$ ) between spectroscopic and petrologic estimates (7 and 0.07 Mt  $\text{SO}_2$ , respectively; BLUTH *et al.*, 1993; DEVINE *et al.*, 1984) by a factor of 40. This gives us to a hope that glaciological estimation will be a useful tool for paleoclimatological studies. Thus, it may be said that  $\text{SO}_2$  emission by past explosive volcanic eruptions in the equatorial region can be estimated by the glaciological method, and that assessment of the impact of past explosive volcanic eruptions on the coeval global climate may be made possible through the study of volcanic signals in glaciers and ice sheets.

## 5. Conclusions

The  $\text{SO}_2$  emission from the 1982 eruption of El Chichon was estimated to be  $17 \pm 1$  Mt  $\text{SO}_2$  by the glaciological method using  $\text{nssSO}_4^{2-}$  concentration in snow samples collected near Dome Fuji Station, Antarctica. This glaciological estimate of  $\text{SO}_2$  emission is consistent with the reported spectroscopic estimate of 7 Mt  $\text{SO}_2$ . This suggests that the glaciological method is applicable to estimate  $\text{SO}_2$  emission from past explosive volcanic eruptions.

## Acknowledgments

We acknowledge Ms. T. KOBAYASHI and Dr. M. IGARASHI of the National Institute of Polar Research for their help in the chemical analysis. We thank Dr. T. FUKUOKA of Rissho University and Mr. N. HAYASHI of Nihon University for their help in the sample preparation. We appreciate the help of the 37th and 38th Japanese Antarctic Research Expeditions, during which snow near Dome Fuji Station was collected.



## References

- ANDRES, R.J., ROSE, W.I., KYLE, P.R., DESILVA, S., FRANCIS, P., GARDEWEG, M. and MORENO ROA, H. (1991): Excessive sulfur dioxide emissions from Chilean volcanoes. *J. Volcanol. Geotherm. Res.*, **46**, 323–329.
- BLUTH, G. J.S., DOIRON, S.D., SCHNETZLER, C.C., KRUEGER, A.J. and WALTER, L.S. (1992): Global tracking of the SO<sub>2</sub> clouds from the June, 1991 Mount Pinatubo eruptions. *Geophys. Res. Lett.*, **19**, 151–154.
- BLUTH, G. J.S., SCHNETZLER, C.C., KRUEGER, A.J. and WALTER, L.S. (1993): The contribution of explosive volcanism to global atmospheric sulphur dioxide concentration. *Nature*, **366**, 327–329.
- CLAUSEN, H.B. and HAMMER, C.U. (1988): The Laki and Tambora eruptions as revealed in Greenland ice cores from 11 locations. *Ann. Glaciol.*, **10**, 16–22.
- DEVINE, J.D., SIGURDSSON, H., DAVIS, A.N. and SELF, S. (1984): Estimates of sulfur and chlorine yield to the atmosphere from volcanic eruptions and potential climatic effects. *J. Geophys. Res.*, **89**, 6309–6325.
- GERLACH, T.M., WESTRICH, H.R., CASADEVALL, T.J. and FINNEGAN, D.L. (1994): Vapor saturation and accumulation in magmas of the 1989–1990 eruption of Redoubt volcano, Alaska. *J. Volcanol. Geotherm. Res.*, **62**, 317–337.
- HAMMER, C.U., CLAUSEN, H.B. and DANSGAARD, D. (1980): Greenland ice sheet evidence of post-glacial volcanism and its climatic impact. *Nature*, **288**, 230–235.
- HATTORI, K. (1996): Occurrence and origin of sulfide and sulfate in the 1991 Mount Pinatubo eruption products. *Fire and Mud: Eruptions and Lahars of Mount Pinatubo, Philippine*, ed. by C.G. NEWHALL and R.S. PUNONGBAYAN. Univ. Washington Press, 807–824.
- IGARASHI, M., KANAMORI, Y. and WATANABE, O. (1998): Analytical method for small amount of polar snow and ice samples by ion chromatography. *Nankyoku Shiryô (Antarct. Rec.)*, **42**, 64–80 (in Japanese with English abstract).
- KAMIYAMA, K., AGETA, Y. and FUJII, Y. (1989): Atmospheric and depositional environments traced from unique chemical compositions of the snow over and inland high plateau, Antarctica. *J. Geophys. Res.*, **94**, 18515–18519.
- KAMIYAMA, K., SHIMADA, W., KITAOKA, K. and EZUMI, S. (1997): Determination of HTO content in polar ice/snow samples by low background liquid scintillation technique. *Nankyoku Shiryô (Antarct. Rec.)*, **41**, 631–642 (in Japanese with English abstract).
- KAZAHAYA, K. and SHINOHARA, H. (1996): Excess degassing of active volcanoes: Processes and mechanisms. *Mem. Geol. Soc. Jpn.*, **46**, 91–104.
- KOHNO, M. (1992): Emission of sulfur, chlorine and fluorine to the atmosphere by volcanic eruptions in Japan in the last 1300 years. M. Sc. thesis, Okayama University, 53 p.
- KOGA, S. and TANAKA, H. (1996): Simulations of seasonal variations of sulfur compounds in the remote marine atmosphere. *J. Atmos. Chem.*, **23**, 163–192.
- KRESS, V. (1997): Magma mixing as a source for Pinatubo sulphur. *Nature*, **389**, 591–593.
- KRUEGER, A. J. (1983): Sighting of El Chichon sulfur dioxide clouds with the Nimbus 7 total ozone mapping spectrometer. *Science*, **220**, 1377–1379.
- KRUEGER, A. J., WALTER, L.S., SCHNETZLER, C.C. and DOIRON, S.D. (1990): TOMS measurement of the sulfur dioxide emitted during the 1985 Nevado del Ruiz eruptions. *J. Volcanol. Geotherm. Res.*, **41**, 7–15.
- LANGWAY, C.C., Jr., CLAUSEN, H.B. and HAMMER, C.U. (1988): An inter-hemispheric volcanic time-marker in ice cores from Greenland and Antarctica. *Ann. Glaciol.*, **10**, 102–108.
- MANDEVILLE, C.W., CAREY, S. and SIGURDSSON, H. (1996): Magma mixing, fractional crystallization and volatile degassing during the 1883 eruption of Krakatau volcano, Indonesia. *J. Volcanol. Geotherm. Res.*, **74**, 243–274.
- MINNIS, P., HARRISON, E.F., STOWE, L.L., GIBSON, G.G., DENN, F.M., DOELLING, D.R. and SMITH, W.L., Jr. (1993): Radiative climate forcing by the Mount Pinatubo eruption. *Science*, **259**, 1411–1415.
- PALAIS, J.M. and SIGURDSSON, H. (1989): Petrologic evidence of volatile emissions from major historic and pre-historic volcanic eruptions. *Understanding Climate Change*, ed. by A. BERGER *et al.* Washington, D. C., Am. Geophys. Union, 31–53 (*Geophys. Monogr.*, 52: IUGG Vol. 7).
- POURCHET, M., PINGLOT, F. and LORIUS, C. (1983): Some meteorological applications of radioactive fallout

- measurements in Antarctic snows. *J. Geophys. Res.*, **88**, 6013–6020.
- POURCHET, M. and PINGLOT, F. (1983): Determination of the stratospheric residence time from the total  $\beta$  activity of Antarctic and Greenland snows. *Geophys. Res. Lett.*, **6**, 365–367.
- RAMPINO, M.R. and SELF, S. (1982): Historic eruptions of Tambora (1815), Krakatau (1883), and Agung (1963), their stratospheric aerosols, and climatic impact. *Quat. Res.*, **18**, 127–143.
- RAMPINO, M.R. and SELF, S. (1984): Sulfur-rich volcanic eruptions and stratospheric aerosols. *Nature*, **310**, 677–679.
- RUTHERFORD, M.J. and DEVINE, J.D. (1996): Preruption pressure-temperature conditions and volatiles in the 1991 dacitic magma of Mount Pinatubo: Source of the giant stratospheric sulfur dioxide cloud. *Fire and Mud: Eruptions and Lahars of Mount Pinatubo, Philippine*, ed. by C.G. NEWHALL and R.S. PUNONGBAYAN. Univ. Washington Press, 751–766.
- SELF, S., ZHAO, J., HOLASEK, R.E., TORRES, R.C. and KING, A.J. (1996): The atmospheric impact of the 1991 Mount Pinatubo. *Fire and Mud: Eruptions and Lahars of Mount Pinatubo, Philippine*, ed. by C.G. NEWHALL and R.S. PUNONGBAYAN. Univ. Washington Press, 1089–1115.
- SIGURDSSON, H. (1982): Volcanic pollution and climate: The 1783 Laki eruption. *EOS*, **63**, no. 32, 601–602.
- SYMONDS, R.B., ROSE, W.I., BLUTH, G.J.S. and GERLACH, T.M. (1994): Volcanic gas studies: Methods, results, and applications. *Rev. Mineral.*, **30**, 1–66.
- UNSCEAR (UNITED NATIONS SCIENTIFIC COMMITTEE ON THE EFFECTS OF ATOMIC RADIATION) REPORT (1982): Ionizing radiation: Sources and biological effects. Report to the General Assembly, with annexes. New York, United Nations. Annex E: 211–248.
- WALLACE, P.J. and GERLACH, T.M. (1994): Magmatic vapor source for sulfur dioxide released during volcanic eruptions: Evidence from Mount Pinatubo. *Science*, **265**, 497–499.
- WATANABE, O., KAMIYAMA, K., MOTOYAMA, H., IGARASHI, M., MATOBA, S. *et al.* (1997): Preliminary report on analyses of melted Dome Fuji ice core obtained in 1993. *Proc. NIPR Symp. Polar Meteorol. Glaciol.*, **11**, 14–23.
- WESTRICH, H.R. and GERLACH, T.M. (1992): Magmatic gas source for the stratospheric SO<sub>2</sub> cloud from the June 15, 1991, eruption of Mount Pinatubo. *Geology*, **20**, 867–870.
- ZIELINSKI, G.A. (1995): Stratospheric loading and optical depth estimates of explosive volcanism over the last 2100 years derived from the Greenland Ice Sheet Project 2 ice core. *J. Geophys. Res.*, **100**, 20937–20955.

(Received May 1, 1998; Revised manuscript accepted June 5, 1998)