

冬季南極海における大気-海氷間の二酸化炭素交換

野村大樹^{1,2,3}, Bruno Delille⁴, Gerhard S. Dieckmann⁵, Jean-Louis Tison⁶,
Klaus M. Meiners^{7,8}, Mats A. Granskog³, 田村岳史⁹

¹ 北大低温研

² 学振

³ ノルウェー極地研

⁴ リエージュ大学

⁵ アルフレッド・ウェグナー極地海洋研

⁶ ベルギー自由大学

⁷ タスマニア大学

⁸ オーストラリア南極局

⁹ 極地研

Sea ice CO₂ flux in the Southern Ocean during mid-winter and early spring

Daiki Nomura^{1,2,3}, Bruno Delille⁴, Gerhard S. Dieckmann⁵, Jean-Louis Tison⁶,
Klaus M. Meiners^{7,8}, Mats A. Granskog³, Takeshi Tamura⁹

¹ *Institute of Low Temperature Science, Hokkaido University*

² *Japan Society for the Promotion of Science*

³ *Norwegian Polar Institute*

⁴ *Université de Liège*

⁵ *Alfred Wegener Institute for Polar and Marine Research*

⁶ *Université Libre de Bruxelles*

⁷ *University of Tasmania*

⁸ *Australian Antarctic Division*

⁹ *National Institute of Polar Research*

There seems little doubt that sea ice is permeable to CO₂ and other gases although air-sea ice gas flux is more or less inhibited at a brine volume fraction of less than 5% representing the threshold for fluid permeability of sea ice. Generally, air-sea ice CO₂ flux is at its minimum in winter due to low sea ice temperatures and consequently reduced permeability despite the fact the partial pressure of CO₂ in sea ice is usually high at that time and sea ice has therefore the potential to release CO₂ to the atmosphere. Here, we present first evidence that snow laden Antarctic sea ice can act as source for atmospheric CO₂ even during mid-winter and early spring. During a mid-winter cruise to the Weddell Sea (AW ECS, 2013) and an early spring cruise off east Antarctica (SIPEX-2, 2012), due to thick insulating snow covers, the bottom of the snow and the surface of the sea ice were relatively warm (>-10°C) even though air temperature was sometimes below -30°C. In addition, in both areas, sea ice was characterized by high bulk-salinities, resulting in brine volume fractions that are generally higher than 5%. Automatic “open-closed” chamber measurements indicated positive CO₂ fluxes of up to +2.5 mmol C m⁻² day⁻¹, illustrating that sea ice acted as a source of atmospheric CO₂. Higher fluxes were measured at bare ice surfaces after removing the snow. However, generally low snow densities (mean: 339 kg m⁻³), indicating a permeable snow cover, facilitated degassing of CO₂ at the snow-air interface. Our results therefore suggest that even in the winter and early spring, Antarctic sea ice can act as CO₂ source for the atmosphere, particularly in areas with a thick insulating snow cover.