

Re-examination of the Himalayan Tethys Sedimentary Sequence

Masaru Yoshida^{1,2} and Bishal Nath Upreti^{1,2}

¹*Gondwana Institute for Geology and Environment, Hashimoto, Japan*

²*Department of Geology, Tri-Chandra Campus, Tribhuvan University, Kathmandu, Nepal*

The central zone of the Himalayan Orogen is occupied by the Tethys Sedimentary Sequence, the Higher Himalayan Gneisses, and the Lesser Himalayan Metasediments juxtaposing with mega faults, being distributed parallel to the mountain range. The lower part of the Tethys Sedimentary Sequence (Neoproterozoic - Cambrian) has been known to be in the unconformity/disconformity relationship with the upper part of the sequence (Ordovician - Eocene) in wide areas along the orogen (e.g., Garzanti et al., 1986; Valdia, 1995; Myrow et al., 2006, 2009). The Tethys Sedimentary Sequence and the Higher Himalayan Gneisses are generally regarded to be juxtaposed with the South Tibetan Detachment fault. However, a gradual relationship including stratigraphic continuity and/or metamorphic gradation has been known from some areas (e.g., Fuchs, 1973/74; Motegi, 2001; Myrow et al., 2003). In relation to the above, Pan-African orogeny in the Himalaya has also been identified by several authors (e.g., Valdia, 1995; Godin, 2003; Gehrels et al., 2006; Yoshida & Upreti, 2007), and the global tectonic set-up of the orogeny has come to be considered (Cawood et al., 2007; Yoshida, 2011).

Based on the evidence and arguments above, the authors propose the terms “Infra Tethys Super Group” and the “Tethys Super Group” to the lower part (Neoproterozoic to Cambro-Ordovician) and the upper part (Middle Ordovician to Eocene) of the so far called Tethys Sedimentary Sequence mentioned above. The protolith of the Higher Himalayan Gneisses is considered to be the Infra Tethys Super Group (e.g., Myrow et al., 2003), taking into account the idea that the gradational relationship between the Infra Tethys Super Group and the Higher Himalayan Gneisses mentioned above represents their original relationship before the detachment. The minimum age of the base of the Infra Tethys Super Group is considered to be the earliest Neoproterozoic (Gehrels et al., 2006).

There are some reports that some formations of the Infra Tethys Super Group are considered to be same as some formations of the Lesser Himalayan Metasediments (Kumar, 1994; Motegi, 2001; Steck, 2003; Myrow et al., 2003; 2006). If so, it is considered that the Infra Tethys Super Group generally conformably overlies the Lesser Himalayan Metasediments in their original structure, before they were separated by the intrusion of the thrust sheet of the Higher Himalayan Gneisses. Taking into account the above idea that the Infra Tethys Super Group is gradational with the Higher Himalayan Gneisses, the idea that the Tethys Super Group, Infra Tethys Super Group, Protolith sedimentary pile of the Higher Himalayan Gneisses, and the Lesser Himalayan Metasediments originally deposited more or less successively along the northern margin of proto Indian subcontinent (Myrow et al., 2003). Shanker et al. (2000) considered that the lower part of the Tethys Sedimentary Sequence (the Infra Tethys Super Group) and the Lesser Himalayan Metasediments formed a part of the Purana Basins sediments formed in the Proto Tethys sea which covered most of the Proto Indian subcontinent and extended northward during the Mesoproterozoic to early Cambrian. It is noted that the above idea is related to the investigation of the protolith of Pan-African high-grade metamorphic rocks of Gondwana.

References cited

- Cawood, P.A., Johnson, M.R.W., Enchain, A.A., 2007, Early Palaeozoic orogenesis along the Indian margin of Gondwana: Tectonic response to Gondwana assembly. *Earth Planet. Sci. Lett.*, 255, 70-84.
- Fuchs. 1973/1974, On the Geology of the Karnali and Dolpo regions, West Nepal. *Mitteilungen der Geologischen Gesellschaft in Wien*, 66-67 Band, 21-32.
- Garzanti, E., Casnedi, R., Jadoul, F., 1986, Sedimentary evidence of a Cambro-Ordovician orogenic event in the northwestern Himalaya. *Sedimentary Geology*, 48, 237-265
- Gehrels, G.E., DeCelles, P.G., Ojha, T.P., Upreti, B.N., 2006, Geologic and U-Th-Pb geochronologic evidence for early Paleozoic tectonism in the Kathmandu thrust sheet, central Nepal Himalaya. *GSA Bull.*, 118, 185-198.
- Godin, L., 2003, Structural evolution of the Tethyan sedimentary sequence in the Annapurna area, central Nepal Himalaya. *J. Asian Earth Sci.*, 22, 307-328.
- Kumar, G. 1997, *Geology of Arunachal Pradesh*. Geol. Soc. India, Bangalore, 217 pages.
- Martin, A.J., Burg, K.D., Kaufman, A.J., Gehrels, G., 2011, Stratigraphic and tectonic implications of field and isotopic constraints on depositional ages of Proterozoic Lesser Himalayan rocks in central Nepal. *Precamb. Res.*, 185, 1-17.
- Myrow, P.M., Hughes, N.C., Paulsen, T.S., Williams, I.S., Pacha, S.K., Thompson, K.R., Bowring, S.A., Peng, S-C.,

- Ahluwalia, A.D., 2003, Integrated tectonostratigraphic analysis of the Himalaya and implications for its tectonic reconstruction. *Earth and Planet. Sci. Let.*, 212, 433-441.
- Myrow, P., Thompson, K.R., Hughes, N.C., Paulsen, T.S., Sell, B.K., Parcha, S.K., 2006, Cambrian stratigraphy and depositional history of the northern Indian Himalaya, Spiti Valley, north-central India. *GSA Bull.*, 118, 491-510.
- Myrow, P., Hughes, N.C., Searle, M.P., Fanning C.M., Peng, S-C, Parcha, S.K., 2009, Stratigraphic correlation of Cambrian Ordovician deposits along the Himalaya: Implications for the age and nature of rocks in the Mount Everest region. *GSA Bull.*, 120, 323-332.
- Motegi, M., 2001, Geology of Bhutan and surrounding areas (1) Outline and problems (in Japanese). Chishitsu Nyusu, Geological Survey of Japan, 567, 6-25.
- Shanker, R., Maithy, P.K., Sing, G., Kumar, G., 2000, Paleogeographic evolution and biotic changes of the Indian subcontinent from the late Palaeoproterozoic to early Palaeozoic. Proceedings volume, International Semi., Precambrian Crust in Easter and Central India, IGCP-368, Bhubaneswar, 34-48.
- Steck, A., 2003, Geology of the NW Indian Himalaya. *Swiss J. Geosciences*, 96, 147-196.
- Valdia, K.S., 1995, Proterozoic sedimentation and Pan-African geodynamic development in the Himalaya. *Precamb. Res.*, 74, 35-55.
- Yoshida, M., 2011, Late Pan-African Peri-Gondwana orogens: Extending the Proto-Himalayan Belt. Presentation at the NIPR Symposium on Geosciences, Nov. 17th, Tachikawa.
- Yoshida, M. & Upreti, B.N., 2007, Commentary on the Pan-African orogeny in the Himalaya. 4th International Symp. on Gondwana to Asia, Fukuoka, IAGR Conf. Ser. 4, 191-193.