

## Horizontal temperature gradients in the polar MLT region (83-105 km) above Tromsø

Satonori Nozawa<sup>1</sup>, Yasunobu Ogawa<sup>2</sup>, H. Fujiwara<sup>3</sup>, Takuo Tsuda<sup>4</sup>, Takuya Kawahara<sup>5</sup>, Norihito Saito<sup>6</sup>, Satoshi Wada<sup>6</sup>,  
Tetsuya Kawabata<sup>1</sup>, Toru Takahashi<sup>2</sup>, Masaki Tsutsumi<sup>2</sup>, Chris Hall<sup>7</sup>, and Asgeir Brekke<sup>8</sup>

<sup>1</sup>ISEE, Nagoya University

<sup>2</sup>NIPR

<sup>3</sup>Faculty of Science and Technology, Seikei University

<sup>4</sup> Department of Computer and Network Engineering, The University of Electro-Communications

<sup>5</sup>Faculty of Engineering, Shinshu University

<sup>6</sup>ASI, RIKEN

<sup>7</sup>TGO, UiT The Arctic University of Norway

<sup>8</sup>Faculty of Science and Technology, UiT The Arctic University of Norway

We have analyzed 2700 hours of temperature data obtained with the Tromsø sodium LIDAR (Nozawa et al. 2014; Kawahara et al. 2017) over 7 seasons between October 2012 and February 2018, and then have calculated horizontal temperature gradients in the polar mesosphere and lower thermosphere (MLT) region between 83 and 105 km. The sodium LIDAR operated at the EISCAT Tromsø site (69.6°N, 19.2°E) has a capability of simultaneous five-directional measurements of temperature and sodium density with good (3 min/500 m) resolutions. Configurations of the sodium LIDAR observational directions are as follows: the vertical position, and south (Azimuth= 180°), north (Azimuth = 0°), west (Azimuth = 270°), and east (Azimuth = 90°) positions. The elevation angle was set to be 77.5° between 2013 and 2016 seasons, while it was 60° for 2 seasons in 2012 and 2017 except for the vertical position. Here we call interval between October and March as “season”, since the sodium LIDAR measurements were made only for the interval (i.e. dark night period).

We have made a statistical study of the temperature gradients. In the statistical study, we have used data sets with their length longer than 4 hours at each night, and then we have used 187 nights in total: winter (i.e. time interval between October 21 and February 23) for 163 nights, and equinox for 24 nights. On average over the 163 nights, the northward temperature gradient is negative (i.e. it was warmer in the south than in the north), and about -0.004 K/km at maximum below 97 km in winter, while it became positive above 97 km. The positive meridional temperature gradient above 97 km is consistent with that of Maeda et al. (2005) who derived temperature gradients utilizing ion temperature data obtained with two EISCAT radars at Longyearbyen (78.2°N, 16.2°E) and Tromsø; the temperature gradients were calculated in a much larger scale (about 800 km) compared to that of this study (about 100 km or less). The averaged zonal temperature gradient was about zero between 85 and 96 km, and it was westward above 96 km. Year-to-year variations are also found: they are more significant above 97 km in the both directions. Southward temperature gradients below 97 km seem to be a common feature over the 6 years except for the 2012 season.

Then, we have investigated variations of temperature gradients on nightly basis. In this talk, we will present results of case studies about the temperature gradients in the polar MLT region. In particular, we focus on effects of auroral activities as well as influence of Sudden Stratospheric Warming (SSW).

### References

- Maeda, S., S. Nozawa, H. Fujiwara, and Y. Ogawa, Comparative study of the high-latitude E-region ion and neutral temperatures in the polar cap and the auroral region derived from the EISCAT radar observations, *J. Geophys. Res.*, *110*, A08301, 10.1029/2004JA010893, 2005.
- Nozawa, S., T. D. Kawahara, N. Saito, C. M. Hall, T. T. Tsuda, T. Kawabata, S. Wada, A. Brekke, T. Takahashi, H. Fujiwara, Y. Ogawa, and R. Fujii, Variations of the neutral temperature and sodium density between 80 and 107 km above Tromsø during the winter of 2010-2011 by a new solid state sodium LIDAR. *J. Geophys. Res.*, *119*, doi:10.1002/2013JA019520, 441-451, 2014.
- Kawahara, T.D., S. Nozawa, N. Saito, T. Kawabata, T.T. Tsuda, and S. Wada, Sodium temperature/wind LIDAR based on laser-diode-pumped Nd:Yag lasers deployed at Tromsø, Norway (69.6°, 19.2°), *Optics Express*, *25*, A491-A501, 2017.