

# SIMULTANEOUS OBSERVATION OF MIDDLE-LEVEL CLOUDS BY A MICROWAVE RADIOMETER AND AN 8.6 mm RADAR (ABSTRACT)

Takao TAKEDA<sup>1</sup>, LIU Guosheng<sup>1</sup> and Makoto WADA<sup>2</sup>

<sup>1</sup>*Water Research Institute, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464*

<sup>2</sup>*National Institute of Polar Research, 9-10, Kaga 1-chome, Itabashi-ku, Tokyo 173*

Simultaneous observation of middle-level clouds was carried out from 1700 to 0800 7 April 1985 by using a 19.35 GHz microwave radiometer and an 8.6 mm vertical pointing radar. The amount of vertically integrated liquid water (ILW) was deduced from the data of microwave radiometer, and radar-echo intensity was converted into the amount of vertically integrated ice water (IIW). Clouds observed can be divided into two different parts. In part 1, the top of the radar-echo was high and air temperature at the top was about  $-30^{\circ}\text{C}$ . Clouds were almost glaciated and the ratio of IIW to total water amount (ILW + IIW) was larger than 50%. It can be said that upper-level clouds which provided a considerable amount of ice particles as seeder clouds existed above middle-level clouds and most of super-cooled water in middle-level clouds was glaciated. On the other hand, in part 2 the top of the radar-echo was lower (air temperature at the top was about  $-15^{\circ}\text{C}$ ) and liquid water was more abundant. The ratio of IIW to total water amount was less than 30%. Clouds were only one layer and sufficient ice particles were not produced in the upper part of middle-level clouds. The amount of vertically integrated liquid water was about  $10\text{ mg/cm}^2$  in part 1 and it exceeded  $40\text{ mg/cm}^2$  in some place of part 2.

(Received April 3, 1986)

# CLOUD STUDIES AT SYOWA STATION IN EAST ANTARCTICA BY MEANS OF LASER-RADAR (ABSTRACT)

Makoto WADA<sup>1</sup> and Yasunobu IWASAKA<sup>2</sup>

<sup>1</sup>*National Institute of Polar Research, 9-10, Kaga 1-chome, Itabashi-ku, Tokyo 173*

<sup>2</sup>*Water Research Institute, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464*

Studies of clouds and precipitation in the polar regions are important for not only cloud physics but also climate research of the world. A lot of studies, mainly on summer clouds in the Arctic region have been already carried out. However, observations for cloud physical approach have been few in the Antarctic region. A laser-radar was set at Syowa Station in East Antarctica since 1983 and clouds in the Antarctic region have been observed by means of the laser-radar. The results of observations are described in the present paper. The wavelength of this ruby laser is  $0.6943\text{ }\mu\text{m}$  and laser power is  $1.0\text{ J/pulse}$ . Seventeen observations of clouds were carried out from May to November in 1983, and the following results have been obtained by analyzing the data. Maximum backscatter coefficient in the clouds was  $1.1\text{ km}^{-1}$ , and this value is larger than the value of backscatter coefficient which was obtained by the observations of clear sky precipitation at South Pole Station, but is much smaller than the value of backscatter coefficient which was obtained by the observations of clouds formed in the middle latitudes. According to a relationship between air temperature and mean depolarization ratio in a cloud, the value of mean depolarization ratio of the cloud