

TERRESTRIAL TARDIGRADES AROUND SYOWA STATION (ABSTRACT)

Kazuo UTSUGI¹ and Yoshikuni OHYAMA²

¹*Department of Biology, Tokyo Women's Medical College,
8-1, Kawadacho, Shinjuku-ku, Tokyo 162*

²*National Institute of Polar Research,
9-10, Kaga 1-chome, Itabshi-ku, Tokyo 173*

Many tardigrades of the Antarctic region have been studied from the beginning of this century. However, in the north-east area where the Japanese Syowa Station is located, only a few species were reported previously.

After these reports, we found five species of tardigrades from the same station and Sør Rondane (K. UTSUGI and Y. OHYAMA: Proc. NIPR Symp. Polar Biol., 2, 190-197, 1989). Later we surveyed Molodezhnaya and Mt. Riiser-Larsen to the east of Syowa Station and found nine species from the samples of both areas (K. UTSUGI and Y. OHYAMA: Proc. NIPR Symp. Polar Biol., 4, 161-170, 1991). The characteristics of these tardigrades were described in these two papers (1989, 1991).

To combine the previous reports and our findings, 10 species of tardigrades including 3 unidentified species are found from Syowa Station (Ongul Islands to Strandnibba). Two species, *Macrobiotus harmsworthi* and *Diphascon conjungens*, are our new findings from the inland area of Sør Rondane. Four species, *Echiniscus kerguelensis*, *Macrobiotus montanus*, *Isohypsibius saracenus* and *Diphascon conjungens* are also our new findings from the coastal areas of Syowa Station, Molodezhnaya and Mt. Riiser-Larsen. In contrast, *M. harmsworthi*, *Hypsibius arcticus* and *Diphascon chilense* are widely distributed in these areas.

It is conceivable that the difference of tardigrade distribution depends on moss and algae flora or the climate in the survey area.

(Received May 2, 1994; Revised manuscript received August 3, 1994)