

Report on Chlorophyll *a* Distribution along the Course of the FUJI in 1981–1982

Yoshio INO* and Mitsuo FUKUCHI**

1981–1982 年「ふじ」航路におけるクロロフィル *a* 量分布報告

伊野良夫*・福地光男**

要旨: 1981–1982 年、第 23 次日本南極地域観測隊の海洋生物定常観測として、「ふじ」船上で行われたクロロフィル *a* 量の測定結果を報告する。

Abstract: Data on chlorophyll *a* concentrations measured on board the FUJI as part of the marine biological programme of the 23rd Japanese Antarctic Research Expedition in 1981–1982 are presented.

Chlorophyll *a* concentrations of surface and subsurface waters were measured on board the icebreaker FUJI as part of the marine biological programme of the 23rd Japanese Antarctic Research Expedition in 1981–1982. Chlorophyll *a* concentration was determined by both the fluorometric and colorimetric methods, using a Shimadzu model RF-500 spectrofluorometer and a HITACHI model 101 spectrophotometer, respectively. A regression of chlorophyll *a* determined by the fluorometric method (*Y*) on chlorophyll *a* determined by the colorimetric method (*X*) was $Y=0.8507X-0.0278$ ($r=0.950$). Data obtained by the fluorometric method are presented in this report.

Surface chlorophyll *a* concentration was measured two to four times a day at 0800, 1300, 1800 and 2300 by the local time along the cruise track of the FUJI between Tokyo and Syowa Station (69°00'S, 39°35'E), Antarctica. A total of 135 stations was occupied between 27 November 1981 and 15 April 1982. The results of measurements are summarized in Table 1 and the surface distribution of chlorophyll *a* is shown in Fig. 1. Subsurface chlorophyll *a* was measured at seven stations in the Indian sector of the Antarctic Ocean between 22 February and 4 March 1982. Water samples from 11 layers down to 200 m depth were collected by the Nansen bottles. The results are listed in Table 2 and the vertical distributions of chlorophyll *a* are shown in Fig. 2.

Acknowledgments

We are indebted to Prof. T. HOSHIAI (National Institute of Polar Research), leader of the JARE-23, and to Captain S. TAKEUCHI of the icebreaker FUJI.

(Received January 13, 1984)

* 早稲田大学教育学部生物学教室. Department of Biology, School of Education, Waseda University, 6-1, Nishi-Waseda 1-chome, Shinjuku-ku, Tokyo 160.

** 国立極地研究所. National Institute of Polar Research, 9-10, Kaga 1-chome, Itabashi-ku, Tokyo 173.




Fig. 1. Surface chlorophyll *a* distribution (—) along the cruise track of the FUJI (----) in 1981-1982. Numeral indicates the serial number of the sampling station. Seven triangles indicate stations at which subsurface chlorophyll *a* concentrations were measured.




Fig. 2. Vertical distribution of chlorophyll *a* at seven stations in the Indian sector of the Antarctic Ocean in February and March 1982 (Stn. 57: 63°02'S, 48°00'E; Stn. 60: 63°00'S, 53°00'E; Stn. 63: 63°00'S, 60°06'E; Stn. 71: 59°26'S, 53°27'E; Stn. 75: 55°16'S, 49°28'E; Stn. 78: 51°04'S, 48°06'E; Stn. 81: 46°40'S, 48°16'E).

Table 1. Data on surface chlorophyll a concentrations and water temperature observed aboard the FUJI between Tokyo and Syowa Station, Antarctica, in 1981–1982.

| Station No. | Date | Time | Latitude | Longitude | Chlorophyll a (mg/m ³) | Water temp. (°C) |
|-------------|---------|------|----------|-----------|---------------------------------------|---------------------|
| 1981 | | | | | | |
| 1 | Nov. 27 | 800 | 26°18'N | 137°47'E | 0.04 | 24.1 |
| 2 | 27 | 1800 | 24 11 | 136 49 | 0.02 | 25.2 |
| 3 | 28 | 800 | 21 13 | 135 15 | 0.03 | 25.9 |
| 4 | 28 | 1800 | 19 23 | 133 55 | 0.03 | 27.1 |
| 5 | 29 | 800 | 16 26 | 132 04 | 0.04 | 28.4 |
| 6 | 29 | 1800 | 14 48 | 130 59 | 0.04 | 28.5 |
| 7 | 30 | 800 | 12 29 | 129 23 | 0.03 | 28.3 |
| 8 | Dec. 1 | 800 | 08 20 | 127 17 | 0.03 | 29.1 |
| 9 | 1 | 1800 | 06 10 | 126 18 | 0.04 | 29.0 |
| 10 | 2 | 800 | 04 00 | 123 51 | 0.09 | 29.0 |
| 11 | 2 | 1800 | 02 33 | 122 18 | 0.27 | 28.8 |
| 12 | 3 | 800 | 01 08 | 120 03 | 0.14 | 28.8 |
| 13 | 3 | 1800 | 00°40'S | 119 13 | 0.12 | 29.4 |
| 14 | 4 | 800 | 03 29 | 118 26 | 0.21 | 29.0 |
| 15 | 4 | 1800 | 05 19 | 117 16 | 0.41 | 28.8 |
| 16 | 5 | 800 | 08 00 | 116 03 | 0.61 | 28.0 |
| 17 | 5 | 1800 | 09 54 | 115 23 | 0.05 | 28.8 |
| 18 | 6 | 800 | 12 57 | 114 43 | 0.06 | 27.8 |
| 19 | 6 | 1800 | 14 44 | 114 19 | 0.06 | 28.3 |
| 20 | 7 | 800 | 17 22 | 113 52 | 0.07 | 26.4 |
| 21 | 7 | 1800 | 19 32 | 113 28 | 0.05 | 25.6 |
| 22 | 8 | 800 | 22 29 | 112 52 | 0.20 | 23.2 |
| 23 | 8 | 1800 | 24 24 | 112 30 | 0.06 | 23.0 |
| 24 | 9 | 800 | 27 06 | 112 45 | 0.08 | 22.2 |
| 25 | 9 | 1800 | 29 07 | 113 25 | 0.11 | 22.8 |
| 26 | 10 | 800 | 31 20 | 114 59 | 0.16 | 20.3 |
| Fremantle | | | | | | |
| 27 | 16 | 1800 | 32 32 | 114 22 | 0.07 | 18.5 |
| 28 | 17 | 815 | 33 59 | 111 36 | 0.09 | 17.7 |
| 29 | 17 | 1800 | 35 00 | 109 46 | 0.21 | 14.4 |
| 30 | 18 | 800 | 37 09 | 108 02 | 0.15 | 15.4 |
| 31 | 18 | 1300 | 38 06 | 108 01 | 0.41 | 13.0 |
| 32 | 18 | 1800 | 39 04 | 108 01 | 0.39 | 12.1 |
| 33 | 18 | 2300 | 40 02 | 107 59 | — | 11.8 |
| 34 | 19 | 800 | 41 47 | 108 00 | 0.37 | 11.4 |
| 35 | 19 | 1300 | 42 46 | 107 59 | 0.34 | 10.7 |
| 36 | 19 | 1800 | 43 45 | 107 59 | 0.30 | 10.2 |
| 37 | 19 | 2300 | 44 38 | 107 43 | 0.31 | 10.1 |
| 38 | 20 | 800 | 46 26 | 107 47 | 0.51 | 9.5 |
| 39 | 20 | 1300 | 47 23 | 107 49 | 0.29 | 6.4 |
| 40 | 20 | 1800 | 48 22 | 107 52 | 0.23 | 6.6 |
| 41 | 20 | 2300 | 49 23 | 107 52 | 0.17 | 4.9 |
| 42 | 21 | 800 | 50 50 | 107 45 | 0.61 | 2.8 |
| 43 | 23 | 640 | 56 49 | 108 00 | 0.18 | 0.6 |
| 44 | 23 | 1730 | 57 41 | 104 50 | 0.23 | 0.1 |
| 45 | 24 | 800 | 58 39 | 99 12 | 0.49 | 0.0 |
| 46 | 24 | 1730 | 59 17 | 95 47 | 0.15 | -0.2 |

Table 1. (continued).

| Station No. | Date | Time | Latitude | Longitude | Chlorophyll <i>a</i> (mg/m ³) | Water temp. (°C) |
|----------------------------|---------|------|----------|-----------|--|---------------------|
| 47 | Dec. 25 | 800 | 60°18'S | 90°21'E | 0.33 | 0.0 |
| 48 | 25 | 1800 | 60 57 | 86 27 | 2.16 | -0.6 |
| 49 | 26 | 800 | 61 50 | 80 50 | 0.66 | -0.6 |
| 50 | 26 | 1800 | 62 27 | 76 58 | 1.34 | -0.4 |
| 51 | 27 | 800 | 63 20 | 71 09 | 1.56 | -0.6 |
| 52 | 27 | 1800 | 63 59 | 66 49 | 0.95 | 0.3 |
| 53 | 28 | 800 | 64 57 | 60 23 | 1.19 | -0.2 |
| 54 | 30 | 800 | 66 16 | 45 11 | 0.18 | -1.6 |
| 55 | 31 | 700 | 66 55 | 40 29 | 0.05 | -1.2 |
| Ice edge off Syowa Station | | | | | | |
| | 1982 | | | | | |
| 56 | Feb. 21 | 1800 | 66 09 | 48 50 | 0.11 | -0.3 |
| 57 | 22 | 930 | 63 02 | 48 00 | 0.13 | 1.1 |
| 58 | 22 | 1800 | 63 00 | 49 35 | 0.18 | 1.2 |
| 59 | 22 | 2300 | 63 00 | 51 12 | 0.18 | 1.1 |
| 60 | 23 | 900 | 63 00 | 53 00 | 0.26 | 1.2 |
| 61 | 23 | 1800 | 62 58 | 56 12 | 0.30 | 0.6 |
| 62 | 23 | 2300 | 62 59 | 57 45 | 0.29 | 1.1 |
| 63 | 24 | 855 | 63 00 | 60 06 | 0.23 | 0.8 |
| 64 | 24 | 1800 | 62 04 | 60 13 | 0.17 | 1.1 |
| 65 | 24 | 2300 | 61 18 | 60 14 | 0.15 | 1.4 |
| 66 | 25 | 800 | 60 00 | 60 01 | 0.23 | 1.4 |
| 67 | 25 | 1800 | 58 55 | 59 59 | 0.13 | 2.1 |
| 68 | 26 | 800 | 59 44 | 60 58 | 0.20 | 1.4 |
| 69 | 26 | 1800 | 59 29 | 58 08 | 0.12 | 2.0 |
| 70 | 26 | 2300 | 59 28 | 56 21 | 0.13 | 2.0 |
| 71 | 27 | 855 | 59 26 | 53 27 | 0.24 | 2.0 |
| 72 | 27 | 1800 | 59 32 | 51 41 | 0.18 | 2.1 |
| 73 | 27 | 2300 | 59 37 | 50 18 | 0.15 | 2.3 |
| 74 | 28 | 800 | 60 17 | 47 59 | 0.17 | 2.0 |
| 75 | Mar. 2 | 855 | 55 16 | 49 28 | 0.07 | 2.9 |
| 76 | 2 | 1800 | 54 09 | 49 28 | 0.08 | 2.7 |
| 77 | 2 | 2300 | 53 04 | 48 27 | 0.07 | 3.6 |
| 78 | 3 | 845 | 51 04 | 48 06 | 0.10 | 4.6 |
| 79 | 3 | 1800 | 49 39 | 48 12 | 0.13 | 5.0 |
| 80 | 3 | 2300 | 48 33 | 48 12 | 0.11 | 6.4 |
| 81 | 4 | 840 | 46 40 | 48 16 | 0.21 | 7.3 |
| 82 | 4 | 1800 | 45 10 | 48 09 | 0.19 | 8.0 |
| 83 | 4 | 2300 | 44 06 | 48 12 | 0.18 | 8.6 |
| 84 | 5 | 800 | 42 29 | 48 17 | 0.55 | 11.9 |
| 85 | 5 | 1300 | 41 44 | 48 35 | 0.33 | 17.8 |
| 86 | 5 | 1800 | 40 42 | 48 53 | 0.16 | 19.7 |
| 87 | 5 | 2300 | 39 35 | 49 03 | 0.13 | 19.2 |
| 88 | 6 | 800 | 37 55 | 49 36 | 0.07 | 19.7 |
| 89 | 6 | 1800 | 35 57 | 50 27 | 0.07 | 21.4 |
| 90 | 7 | 800 | 33 05 | 51 39 | 0.05 | 22.7 |
| 91 | 7 | 1800 | 31 04 | 52 16 | 0.07 | 24.5 |
| 92 | 8 | 800 | 28 37 | 53 14 | 0.15 | 25.0 |
| 93 | 8 | 1800 | 26 49 | 54 09 | 0.05 | 26.2 |

Table 1. (continued).

| Station No. | Date | Time | Latitude | Longitude | Chlorophyll α (mg/m ³) | Water temp. (°C) |
|-------------|--------|------|----------|-----------|--|---------------------|
| 94 | Mar. 9 | 800 | 24°11'S | 55°20'E | 0.05 | 26.8 |
| 95 | 9 | 1800 | 22 14 | 55 55 | 0.06 | 28.2 |
| Port Louis | | | | | | |
| 96 | 17 | 1800 | 19 19 | 58 31 | 0.04 | 27.2 |
| 97 | 18 | 800 | 18 00 | 60 42 | 0.01 | 27.6 |
| 98 | 18 | 1800 | 17 03 | 62 14 | 0.02 | 28.4 |
| 99 | 19 | 800 | 15 42 | 64 18 | 0.02 | 27.8 |
| 100 | 19 | 1800 | 14 45 | 66 05 | 0.02 | 27.7 |
| 101 | 20 | 800 | 12 57 | 68 29 | 0.05 | 28.0 |
| 102 | 20 | 1800 | 11 50 | 70 15 | 0.05 | 28.6 |
| 103 | 21 | 800 | 10 16 | 72 38 | 0.04 | 28.4 |
| 104 | 21 | 1800 | 09 05 | 74 24 | 0.03 | 28.8 |
| 105 | 22 | 800 | 07 19 | 76 56 | 0.04 | 28.8 |
| 106 | 22 | 1800 | 06 02 | 78 28 | 0.05 | 29.1 |
| 107 | 23 | 800 | 04 21 | 80 43 | 0.04 | 28.5 |
| 108 | 23 | 1800 | 03 27 | 81 49 | 0.03 | 28.9 |
| 109 | 24 | 800 | 01 41 | 84 06 | 0.04 | 28.8 |
| 110 | 24 | 1800 | 00 28 | 85 44 | 0.05 | 30.2 |
| 111 | 25 | 800 | 01°06'N | 87 59 | 0.04 | 28.9 |
| 112 | 25 | 1800 | 02 11 | 89 42 | 0.04 | 29.3 |
| 113 | 26 | 800 | 03 48 | 91 55 | 0.04 | 28.9 |
| 114 | 26 | 1800 | 05 01 | 93 29 | 0.04 | 30.0 |
| 115 | 27 | 800 | 06 10 | 95 18 | 0.12 | 29.4 |
| 116 | 27 | 1800 | 05 45 | 96 46 | 0.11 | 29.7 |
| 117 | 28 | 800 | 04 52 | 98 33 | 0.13 | 29.1 |
| 118 | 28 | 1800 | 04 07 | 99 48 | 0.13 | 29.7 |
| 119 | 30 | 800 | 03 04 | 100 34 | 0.31 | 29.4 |
| 120 | 30 | 1800 | 02 07 | 102 05 | 0.55 | 29.1 |
| 121 | 31 | 800 | 01 11 | 103 50 | 0.45 | 28.9 |
| Singapore | | | | | | |
| 122 | Apr. 8 | 1800 | 01 43 | 105 05 | 0.12 | 29.6 |
| 123 | 9 | 800 | 02 40 | 107 37 | 0.09 | 29.4 |
| 124 | 9 | 1800 | 04 11 | 108 57 | 0.05 | 29.2 |
| 125 | 10 | 800 | 06 37 | 110 14 | 0.06 | 28.6 |
| 126 | 10 | 1800 | 08 05 | 111 00 | 0.04 | 28.4 |
| 127 | 11 | 800 | 10 29 | 112 36 | 0.05 | 28.1 |
| 128 | 11 | 1800 | 12 06 | 113 46 | 0.04 | 27.6 |
| 129 | 12 | 800 | 14 19 | 115 14 | 0.06 | 28.2 |
| 130 | 12 | 1800 | 16 09 | 116 06 | 0.04 | 26.3 |
| 131 | 13 | 800 | 18 33 | 118 03 | 0.04 | 25.0 |
| 132 | 13 | 1800 | 19 34 | 120 16 | 1.12 | 24.7 |
| 133 | 14 | 800 | 21 10 | 123 10 | 0.05 | 23.9 |
| 134 | 14 | 1800 | 22 48 | 124 54 | 0.06 | 23.4 |
| 135 | 15 | 800 | 24 49 | 127 03 | 0.07 | 22.0 |

Table 2. Data on subsurface chlorophyll *a*, phaeophytin and pigment ratio observed at seven stations in the Indian sector of the Antarctic Ocean in February and March 1982.

| Station No. | Date Time | Position | Depth (m) | Chl. <i>a</i> (mg/m ³) | Phaeopigments (mg/m ³) | Pigment ratio (%) |
|-------------|----------------------|--------------------|-----------|------------------------------------|------------------------------------|-------------------|
| 57 | 1982 Feb. 22 0930 | 63°02'S 48°00'E | 0 | 0.13 | 0.07 | 65.6 |
| | | | 10 | 0.16 | 0.07 | 69.0 |
| | | | 25 | 0.09 | 0.13 | 42.1 |
| | | | 39 | 0.12 | 0.09 | 56.7 |
| | | | 59 | 0.11 | 0.03 | 76.1 |
| | | | 79 | 0.15 | 0.15 | 50.1 |
| | | | 98 | 0.22 | 0.21 | 51.0 |
| | | | 123 | 0.08 | 0.12 | 40.6 |
| | | | 148 | 0.02 | 0.01 | 25.0 |
| | | | 172 | 0.01 | 0.06 | 9.7 |
| | | | 197 | 0.00 | 0.09 | 1.8 |
| 60 | 1982 Feb. 23 0900 | 63°00'S 53°00'E | 0 | 0.26 | 0.14 | 65.5 |
| | | | 10 | 0.31 | 0.12 | 71.7 |
| | | | 24 | 0.23 | 0.12 | 65.5 |
| | | | 39 | 0.28 | 0.07 | 79.2 |
| | | | 58 | 0.30 | 0.01 | 97.0 |
| | | | 78 | 0.16 | 0.08 | 64.8 |
| | | | 97 | 0.18 | 0.12 | 59.3 |
| | | | 122 | 0.10 | 0.11 | 46.7 |
| | | | 146 | 0.05 | 0.06 | 43.7 |
| | | | 171 | 0.03 | -0.01 | — |
| | | | 195 | 0.00 | 0.06 | 7.3 |
| | | | 0 | 0.23 | 0.16 | 58.9 |
| 63 | 1982 Feb. 24 0855 | 63°00'S 60°06'E | 10 | 0.31 | 0.56 | 35.6 |
| | | | 24 | 0.32 | 0.45 | 41.6 |
| | | | 39 | 0.30 | 0.48 | 37.9 |
| | | | 58 | 0.25 | 0.38 | 40.2 |
| | | | 77 | 0.21 | 0.32 | 39.2 |
| | | | 97 | 0.29 | 0.47 | 38.0 |
| | | | 121 | 0.21 | 0.34 | 38.2 |
| | | | 145 | 0.06 | 0.07 | 46.1 |
| | | | 169 | 0.02 | 0.07 | 21.8 |
| | | | 193 | 0.01 | 0.06 | 12.1 |
| | | | 0 | 0.24 | 0.09 | 72.3 |
| 71 | 1982 Feb. 27 0855 | 59°26'S 53°27'E | 10 | 0.30 | 0.05 | 84.6 |
| | | | 24 | 0.24 | 0.06 | 79.0 |
| | | | 39 | 0.23 | 0.10 | 70.4 |
| | | | 58 | 0.22 | 0.11 | 67.4 |
| | | | 78 | 0.11 | 0.11 | 49.5 |
| | | | 97 | 0.06 | 0.06 | 50.3 |
| | | | 121 | 0.03 | 0.08 | 29.1 |
| | | | 146 | 0.02 | 0.01 | 72.8 |
| | | | 170 | 0.01 | 0.01 | 58.2 |
| | | | 194 | 0.00 | 0.01 | 43.7 |

Table 2. (continued).

| Station No. | Date Time | Position | Depth (m) | Chl. α (mg/m 3) | Phaeopigments (mg/m 3) | Pigment ratio (%) |
|-------------|---------------------|--------------------|-----------|----------------------------|----------------------------|-------------------|
| 75 | 1982 Mar. 2 0855 | 55°16'S 49°28'E | 0 | 0.07 | 0.02 | 76.3 |
| | | | 9 | 0.08 | 0.03 | 69.9 |
| | | | 23 | 0.06 | 0.04 | 58.2 |
| | | | 38 | 0.04 | 0.06 | 43.7 |
| | | | 56 | 0.05 | 0.05 | 48.5 |
| | | | 75 | 0.07 | 0.04 | 61.1 |
| | | | 94 | 0.07 | 0.04 | 62.6 |
| | | | 117 | 0.05 | 0.02 | 72.8 |
| | | | 141 | 0.03 | 0.02 | 58.2 |
| | | | 164 | 0.01 | 0.05 | 17.5 |
| 78 | 1982 Mar. 3 0845 | 51°04'S 48°06'E | 0 | 0.10 | 0.05 | 69.1 |
| | | | 10 | 0.12 | 0.06 | 65.5 |
| | | | 25 | 0.14 | 0.08 | 61.9 |
| | | | 40 | 0.10 | 0.09 | 53.9 |
| | | | 59 | 0.12 | 0.05 | 69.1 |
| | | | 79 | 0.12 | 0.00 | 96.6 |
| | | | 99 | 0.10 | 0.02 | 80.7 |
| | | | 124 | 0.07 | 0.13 | 34.8 |
| | | | 149 | 0.04 | 0.03 | 55.8 |
| | | | 173 | 0.02 | 0.04 | 34.0 |
| 81 | 1982 Mar. 4 0840 | 46°40'S 48°16'E | 0 | 0.21 | 0.12 | 63.5 |
| | | | 9 | 0.20 | 0.12 | 61.7 |
| | | | 22 | 0.16 | 0.06 | 72.8 |
| | | | 35 | 0.15 | 0.07 | 67.0 |
| | | | 53 | 0.12 | 0.08 | 59.8 |
| | | | 71 | 0.15 | 0.07 | 69.9 |
| | | | 88 | 0.11 | 0.12 | 47.1 |
| | | | 110 | 0.08 | 0.15 | 34.0 |
| | | | 132 | 0.05 | 0.13 | 27.3 |
| | | | 155 | 0.03 | 0.10 | 24.3 |
| | | | 177 | 0.01 | 0.03 | 29.1 |